










Reproductive genetics

A genetic variant in the 3'-UTR of *PIWIL4* confers risk for extreme phenotypes of male infertility by altering miR-215 and miR-136 binding affinity

Sara González-Muñoz^{1,2,†}, Miriam Cerván-Martín¹ ^{3,†}, Andrea Guzmán-Jiménez^{1,2}, Ana Isabel Rodríguez-Martín¹, Nicolás Garrido⁴, José A. Castilla^{2,5}, M. Carmen Gonzalvo^{2,6}, Ana Clavero^{2,6}, Marta Molina¹ ^{2,6}, Miguel Ángel Vilches⁷, Andrea Espuch-Oliver⁸, Vicente Maldonado⁹, María Luisa García-Peña⁹, Noelia Galiano-Gutiérrez¹⁰, Esther Santamaría¹⁰, Cristina González⁴, Fernando Quintana-Ferraz¹ ⁴, Susana Gómez⁴, David Amorós⁴, Luis Martínez-Granados¹¹, Yanira Ortega-González¹ ¹², Miguel Burgos¹, Iris Pereira-Caetano¹³, Graça S. Pinto¹⁴, Ana Aguiar¹⁵, Isabel S. Pereira^{15,16}, Olga López-Rodrigo¹⁷, Lluís Bassas¹⁷, Susana Seixas¹ ^{18,19}, João Gonçalves^{13,20}, Alexandra M. Lopes^{19,21}, Sara Larriba¹ ²², Lara Bossini-Castillo¹ ^{1,2}, F. David Carmona¹ ^{1,2,*†}, and Rogelio J. Palomino-Morales¹ ^{2,23,*†}

¹Departamento de Genética e Instituto de Biotecnología, Centro de Investigación Biomédica (CIBM), Universidad de Granada, Granada, Spain

²Instituto de Investigación Biosanitaria, ibs.GRANADA, Granada, Spain

³Institute of Parasitology and Biomedicine López-Neyra (IPBLN), CSIC, Granada, Spain

⁴IVI Foundation, Health Research Institute La Fe, Valencia, Spain

⁵Departamento de Anatomía y Embriología Humana, Universidad de Granada, Granada, Spain

⁶Unidad de Reproducción, UGC Obstetricia y Ginecología, HU Virgen de las Nieves, Granada, Spain

⁷Ovoclinic & Ovobank, Clínicas de Reproducción Asistida y Banco de óvulos, Marbella, Málaga, Spain

⁸Unidad de Reproducción Humana Asistida, Hospital Universitario Torrecárdenas, Almería, Spain

⁹UGC de Obstetricia y Ginecología, Complejo Hospitalario de Jaén, Jaén, Spain

¹⁰Vida Sevilla, Vida Medicina Reproductiva, Sevilla, Spain

¹¹Hospital Universitario Príncipe de Asturias, Alcalá de Henares, Madrid, Spain

¹²Unidad de Urología, Hospital Universitario de Canarias, Santa Cruz de Tenerife, Spain

¹³Departamento de Genética Humana, Instituto Nacional de Saúde Dr. Ricardo Jorge, Lisbon, Portugal

¹⁴Centro de Medicina Reprodutiva, Maternidade Dr Alfredo da Costa, Unidade Local de Saúde São José, Lisboa, Portugal

¹⁵Unidade de Medicina da Reprodução, Departamento de Obstetrícia, Ginecologia e Medicina da Reprodução, Unidade Local de Saúde de Santa Maria, Lisboa, Portugal

¹⁶Centro de Medicina da Reprodução, Hospital Central do Funchal Dr. Nélio Mendonça, Funchal, Portugal

¹⁷Laboratory of Andrology and Sperm Bank, Andrology Service-Fundació Puigvert, Barcelona, Spain

¹⁸IS—Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Porto, Portugal



¹⁹Institute of Molecular Pathology and Immunology, University of Porto (IPATIMUP), Porto, Portugal

²⁰Comprehensive Health Research Centre, Nova Medical School, Lisbon, Portugal

²¹CGPP-IBMC—Centro de Genética Preditiva e Preventiva, Instituto de Biologia Molecular e Celular, Universidade do Porto, Porto, Portugal

²²Immune-Inflammatory Processes and Gene Therapeutics Group; Genes, Disease and Therapy Program; Bellvitge Biomedical Research Institute (IDIBELL), L'Hospitalet de Llobregat, Barcelona, Spain

²³Departamento de Bioquímica y Biología Molecular I, Centro de Investigación Biomédica (CIBM), Universidad de Granada, Granada, Spain

*Correspondence address. Centro de Investigación Biomédica (CIBM), Universidad de Granada, Parque Tecnológico Ciencias de la Salud, Avda. del Conocimiento, S/N. 18016-Armilla (Granada), Spain. Tel: +34-958243088. E-mail: rpm@ugr.es  <https://orcid.org/0000-0003-3507-343X> (R.J.P.-M.); Centro de Investigación Biomédica (CIBM), Universidad de Granada, Parque Tecnológico Ciencias de la Salud, Avda. del Conocimiento, S/N. 18016-Armilla (Granada), Spain. Tel: +34-958241543. E-mail: dcarmona@ugr.es  <https://orcid.org/0000-0002-1427-7639> (F.D.C.)

[†]These authors contributed equally to this work.

[‡]These authors shared senior authorship.

ABSTRACT

STUDY QUESTION: What is the functional impact of the rs508485 genetic polymorphism, located in the 3'-untranslated region (UTR) region of the *PIWIL4* gene, on non-obstructive azoospermia (NOA)?

SUMMARY ANSWER: The rs508485 genetic variant contributes to the pathogenesis of extreme patterns of NOA by modulating *PIWIL4* expression through microRNA (miRNA) interactions.

Received: December 23, 2024. Revised: April 23, 2025. Editorial decision: May 15, 2025.

© The Author(s) 2025. Published by Oxford University Press on behalf of European Society of Human Reproduction and Embryology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com.

WHAT IS KNOWN ALREADY: Male infertility represents a significant global health challenge with profound societal and economic consequences. One of the most severe forms of male infertility is NOA, which is characterized by severe spermatogenic failure (SPGF) of idiopathic origin in most cases. Cumulating knowledge increasingly suggests that this idiopathic form of NOA may represent a multifactorial condition involving complex interactions between genetic and environmental factors. The PIWI protein subfamily, particularly PIWIL4, plays a pivotal role in spermatogenesis by processing PIWI-interacting RNAs, which silence retrotransposons to protect genomic integrity. Genetic variations in this gene have been found to be associated with susceptibility to NOA.

STUDY DESIGN, SIZE, DURATION: A case–control study was conducted in a European cohort including 1516 infertile men with SPGF and 2451 fertile controls. Logistic regression and functional assays were employed to investigate the functional role of the rs508485 polymorphism in *PIWIL4*.

PARTICIPANTS/MATERIALS, SETTING, METHODS: Participants were genotyped for the rs508485 polymorphism. Associations between the polymorphism and NOA phenotypes, including Sertoli cell-only (SCO) syndrome and testicular sperm extraction (TESE) outcomes, were assessed. *In silico* tools predicted miRNA binding effects, which were subsequently validated using luciferase reporter assays.

MAIN RESULTS AND THE ROLE OF CHANCE: The T allele of rs508485 was significantly associated with the SCO phenotype ($P = 2.69E-03$, $OR = 1.34$) and unfavourable TESE outcomes ($P = 1.09E-03$, $OR = 1.54$). *In silico* analyses predicted that the rs508485 variant might alter binding sites in the 3'-UTR region of *PIWIL4* for different miRNAs, such as hsa-miR-215-3p and hsa-miR-136-3p. Functional validation using luciferase assays confirmed that these miRNAs differentially bind to the T and C alleles of this polymorphism, influencing *PIWIL4* regulation.

LARGE SCALE DATA: N/A.

LIMITATIONS, REASONS FOR CAUTION: The study is limited to a single genetic polymorphism and functional assays were performed *in vitro*. Additional studies are required to validate these findings across diverse populations and explore additional genetic interactions.

WIDER IMPLICATIONS OF THE FINDINGS: These findings highlight the critical role of miRNA regulation in extreme forms of male infertility by influencing the expression of essential spermatogenesis genes, such as *PIWIL4*. Our study sheds light on the genetic mechanisms underlying spermatogenesis and suggests potential therapeutic targets for NOA.

STUDY FUNDING/COMPETING INTEREST(S): This work was supported by the Spanish Ministry of Economy and Competitiveness through the Spanish National Plan for Scientific and Technical Research and Innovation (grant: PID2020-120157RB-I00 funded by MICIU/AEI/10.13039/501100011033; and grant: PID2023-152215OB-I00 funded by MICIU/AEI/10.13039/501100011033 and ERDF, EU). M. C.-M is beneficiary of a Juan de la Cierva 2022 postdoctoral fellowship (reference number JDC2022-048561-I) funded by MCIN/AEI/10.13039/501100011033 and European Union NextGenerationEU/PRTR. S.L. received support from “Generalitat de Catalunya” (grant 2021SGR052). S.L. is sponsored by the “Researchers Consolidation Program” from the SNS-Dpt. Salut Generalitat de Catalunya (Exp. CES09/020). S.G.-M. was funded by grant ref. FPU23/02674. S.S. was supported by funds from the Portuguese Foundation for Science and Technology (FCT; 10.54499/DL57/2016/CP1363/CT0019). The authors declare no competing interests.

Keywords: male infertility / NOA / SCO / TESE / *PIWIL4* / piRNA / miRNA / SNP

Introduction

Infertility is currently recognized as a significant global public health concern by the World Health Organization (WHO). This condition affects ~10–15% of couples worldwide, with male factor infertility accounting for around half of these cases (Massart et al., 2012; Tournaye et al., 2017; World Health Organization, 2017). In this regard, non-obstructive azoospermia (NOA), defined as the total absence of sperm in the ejaculate due to a severe spermatogenic failure (SPGF), represents the most extreme and one of the most common causes of male infertility, affecting 10% of infertile men and 1% of the male population (Cocuzza et al., 2013; Tournaye et al., 2017). While several Mendelian causes have been identified for NOA, the aetiology of this condition remains unknown for around 75% of affected men. Increasing evidence clearly suggests that this idiopathic form of NOA represents a complex trait influenced by interactions between genetic and environmental factors (Cervan-Martin et al., 2020, 2022, 2024).

P-element-induced wimpy testis (PIWI) proteins, part of the Argonaut protein subfamily, include PIWIL1, PIWIL2, PIWIL3, and PIWIL4 in humans. These proteins are predominantly expressed in germline cells and play a critical role in silencing retrotransposons through binding to PIWI-interacting RNAs (piRNAs), a class of small non-coding RNAs (Aravin et al., 2007a; Ozata et al., 2019). This silencing is achieved via mechanisms such as DNA methylation and histone modification, both of which are essential for spermatogenesis (Hartig et al., 2007; Klattenhoff and Theurkauf, 2008; Juliano et al., 2011). Although piRNA expression has been detected in both testis and ovaries, only male knock-out mice for

orthologous *PIWI* genes develop infertility, likely due to upregulation of transposable elements in the germline (Aravin et al., 2007b; He et al., 2009). Specifically, several studies have revealed that *PIWIL4* deficiency in male mice leads to meiotic arrest (MA) during spermatogenesis, resulting in male sterility (Deng and Lin, 2002; Kuramochi-Miyagawa et al., 2004; Carmell et al., 2007). Regarding common genetic variations in the *PIWIL4* gene, several studies have reported associations of different single-nucleotide polymorphisms (SNPs), including rs508485, with a higher risk of male infertility due to SPGF in populations of different ancestries (Gu et al., 2010; Munoz et al., 2014; Kamaliyan et al., 2017, 2018). However, the functional implications of the two allelic variants of rs508485 in the pathophysiology of male infertility remain unclear. There is ongoing debate regarding its association with specific patterns of SPGF, such as NOA, severe oligozoospermia (SO, fewer than 5 million spermatozoa per millilitre of ejaculate), or even a subgroup of NOA patients with a histological phenotype of MA (Gu et al., 2010; Munoz et al., 2014; Kamaliyan et al., 2017, 2018). Although some authors have hypothesized that the rs508485 variant affects mRNA stability or alters the binding affinity of regulatory microRNAs (miRNAs), the precise molecular mechanism in which this SNP exerts its influence remains ambiguous (Kamaliyan et al., 2018).

Based on the considerations exposed above, we conducted the present study to investigate the possible association between the *PIWIL4*-rs508485 SNP and NOA in a well-powered and characterized cohort of European descent. We aimed to elucidate the functional implications of this variant in the pathogenesis of this form of male infertility.

Materials and methods

Study population

Two independent case-control populations of European descent were analysed in this study, comprising a total of 1516 infertile men due to SPGF and 2451 unaffected controls. The discovery cohort included Iberian and German individuals (1274 cases and 1951 controls) and the replication cohort was composed only of Iberian men (242 cases and 500 controls).

All participants provided informed written consent and DNA samples were irreversibly anonymized prior to enrolment. The study procedures adhered to the principles outlined in the Declaration of Helsinki and received approval from the institutional Ethics Committee 'CEIM/CEI Provincial de Granada' (Andalusia, Spain), ensuring full compliance with government policies and ethical standards.

The discovery cohort data utilized in this study were derived from a previously published genome-wide association study (GWAS), where detailed clinical and demographic information was provided (Cervan-Martin *et al.*, 2022). The same inclusion criteria were applied for the replication cohort. Briefly, this study included men diagnosed with SPGF of idiopathic origin based on WHO guidelines (World Health Organization, 2021). Diagnosis was established through comprehensive clinical assessment, including physical examination, endocrine profiling, and histological evaluation when available, to confirm the total absence of sperm in the ejaculate. To ensure a well-defined NOA cohort, known causes of infertility were systematically excluded, including anatomical disorders, testicular neoplasms, obstructive cases, congenital bilateral absence of vas deferens, as well as genetic abnormalities such as karyotype anomalies and Y chromosome microdeletions, both of which were determined through standard genetic screening. Around half of the NOA cohort underwent a testicular sperm extraction (TESE) procedure from a testis biopsy, allowing classification into three different histological subgroups: hypospermatogenesis (HS, production of an extremely low number of sperm cells), MA (incomplete differentiation of the germline), and Sertoli cell-only (SCO) (total absence of germ cells in the seminiferous tubules). Moreover, NOA patients were also divided into two additional subgroups based on the TESE outcome (i.e. successful and unsuccessful TESE). The control group consisted of fertile men, matched with cases based on geographical origin and average age, who had fathered one or more healthy children naturally. [Supplementary Table S1](#) presents comprehensive details regarding the study cohort.

Genotyping, quality controls, and imputation

DNA samples from the replication cohort were extracted from peripheral blood mononuclear cells and genotyped using the Infinium Global Screening Array-24 v3.0 (GSA) on an iScan System (Illumina, Inc). This procedure followed the same protocol outlined for the samples from the discovery cohort, as detailed in our previous study (Cervan-Martin *et al.*, 2022).

The SNP rs508485, selected from the discovery cohort using GWAS data (Cervan-Martin *et al.*, 2022), was genotyped in the replication cohort using the TaqManTM allelic discrimination technology on a 7900HT Fast Real-Time PCR System with a pre-designed TaqMan probe (reference assay ID: C_1365626_10). The SDS 2.3 software was used for allelic discrimination (Thermo Fisher ScientificTM).

Statistical analysis

Logistic regression models were used to compare all case groups (SPGF, NOA, MA, HS, SCO, and TESEneg) against unaffected

controls, incorporating covariates such as principal components and country of origin and assuming additive, dominant and recessive effects. Additionally, a combined analysis of both cohorts was performed via inverse variance-weighted meta-analysis under a fixed-effects model. The possible heterogeneity of the estimated odds ratios (OR) between studies was evaluated using I^2 and Cochran's Q tests. The P-value for statistical significance was set at $P=0.05$.

Functional relevance of the rs508485 polymorphism

To assess the functional impact of PIWIL4-rs508485 on the spermatogenic process, functional annotations of the human genome overlapping the SNP under study (rs508485) and all its proxies (polymorphisms within the same linkage disequilibrium (LD) block, $R^2 \geq 0.8$) were retrieved from public databases. Initially, we identified proxies for the associated rs508485 variant in the European population using LDlink (Machiela and Chanock, 2015). To enhance our understanding of the regulatory potential of the selected variants, we extracted detailed functional annotation data from multiple publicly available resources including HaploReg v.4.2. (Ward and Kellis, 2016), RegulomeDB (Dong and Boyle, 2019), Open Target Genetics Portal (Ghousssaini *et al.*, 2021), GTEx (v8) (GTEx Consortium, 2020), and SNP Nexus (Oscano *et al.*, 2020).

Prediction of miRNA target sequences

A comprehensive bioinformatic analysis was performed to predict the target sequences of miRNAs that were altered by the rs508485 risk allele located in the 3'-untranslated region (UTR) region of PIWIL4. For this purpose, the following bioinformatics tools were utilized: TargetScan (McGeary *et al.*, 2019), PolymiRTS (Bhattacharya *et al.*, 2014), miRBase (Kozomara *et al.*, 2019), and miRmap (Vejnar and Zdobnov, 2012), which assess the binding potential based on sequence complementarity and seed region match. Only candidate target sites identified by at least two prediction algorithms were selected for further analysis. Additionally, RNAhybrid (Kruger and Rehmsmeier, 2006) was employed to determine the minimum free energy (MFE) required for the hybridization of miRNAs to their predicted target sites (lower MFE values indicate enhanced stability of formed structures).

Cell cultures

HEK-293T cells (obtained from ATCC[®] and provided by Dr José Luis Garcia-Perez, GENYO, Spain) were cultured in Dulbecco's modified Eagle's medium supplemented with 10% foetal bovine serum (Gibco), 100 U/ml penicillin G, and 100 µg/ml streptomycin (Invitrogen). Cell cultures were maintained in humidified incubators at 37°C with 5% CO₂. Mycoplasma contamination was ruled out monthly using a PCR-based assay (Minerva).

Plasmid constructs

In order to enhance the possible effect of the PIWIL4-rs508485 risk variant, five copies of the 3'-UTR region of PIWIL4 were cloned into the psiCHECK2 vector. Specifically, 56 bp segments of this region were selected to minimize secondary structures, based on predictions generated by the RNAfold tool (Lorenz *et al.*, 2011). These segments, containing either the T or C allele of the rs508485 polymorphism, along with a mutated miRNA binding sequence as a negative control are described in [Supplementary Table S2](#). All constructs were synthesized in pUC57 (GenScript) and cloned into psiCHECK2 using XhoI and NotI restriction enzyme sites (New England Biolabs). Correct insertions were confirmed by restriction digestion and Sanger sequencing.

miRNAs mimics

miRIDIAN miRNA mimics for hsa-miR-215-3p, hsa-miR-136-3p, and a scrambled control (Supplementary Table S3) were purchased from Dharmacon. Mimics were resuspended in 1× siRNA Buffer (Fisher Scientific) to a working concentration of 20 μM and stored at −80°C until use.

Transfection and luciferase reporter assays

HEK-293T cells were seeded (1×10^5 per well in 24-well plates) 24 h prior to transfection. Cells at 80–90% confluence were transfected with 10 ng of each psiCHECK2 plasmid and varying concentrations (10, 25, and 50 nM) of selected miRNA mimics using DharmafECT reagents (Dharmacon).

Dual luciferase assays were conducted to confirm specific miRNA binding using the Dual Luciferase Reporter Assay System (Promega) and GloMax[®] 96 Microplate Luminometer (Promega), following manufacturer's instructions. Three experiments per condition with two measurements per sample were performed. Luciferase activity was measured as relative luminescence units (renilla/firefly luciferase) and normalized to scrambled controls. Statistical significance was determined using a two-tailed Student's t-test.

Results

Testing for association of rs508485 with male infertility due to SPGF

We initially compared the allele frequencies of the rs508485 SNP between the overall SPGF group (encompassing both NOA and SO patients) and the control group to assess its role in SPGF susceptibility. No statistically significant differences were found across genetic models (Table 1, Supplementary Table S4). Subsequently, and taking advantage of the detailed patient phenotyping, we compared the NOA group as well as the different NOA subgroups, defined by specific histological patterns (i.e. HS, MA, and SCO) and unsuccessful TESE outcome (TESEneg), against the control group. Notably, *PIWIL4*-rs508485 was associated with both TESEneg and SCO, showing significant *P*-values for the recessive and additive models, respectively, in the combined European discovery cohort of Iberians ($P_{\text{TESEneg, rec}}=3.01\text{E-}02$, OR (CI) $_{\text{TESEneg, rec}}=1.54$ (1.04–2.28); $P_{\text{SCO, add}}=4.10\text{E-}02$, OR (CI) $_{\text{SCO, add}}=1.35$ (1.01–1.81)) and Germans ($P_{\text{TESEneg, rec}}=1.03\text{E-}02$, OR (CI) $_{\text{TESEneg, rec}}=1.70$ (1.13–2.54); $P_{\text{SCO, add}}=3.82\text{E-}02$, OR (CI) $_{\text{SCO, add}}=1.36$ (1.02–1.81)) (Table 1, Supplementary Table S4).

To validate these findings, we conducted an additional case-control analysis in an independent Iberian cohort. While this replication analysis did not yield significant *P*-values due to smaller subgroup sizes, effect sizes similar to the discovery cohort were observed for the SCO versus control and TESEneg versus control comparisons (OR = 1.25 and OR = 1.17, respectively) (Table 1, Supplementary Table S4). Interestingly, the meta-analysis combining both cohorts substantially increased the statistical significance observed in the discovery cohort for SCO ($P_{\text{META}}=2.69\text{E-}03$, OR $_{\text{META}}=1.34$, CI $_{\text{META}}=1.11$ –1.63) and TESEneg ($P_{\text{META}}=1.09\text{E-}03$, OR $_{\text{META}}=1.54$, CI $_{\text{META}}=1.19$ –2.00) conditions, with no significant heterogeneity observed between studies (Table 1, Supplementary Table S4).

In silico characterization of the rs508485 polymorphism and its proxies

With the aim of investigating the biological impact of the rs508485 SNP in SCO development and TESE outcome, we analysed the functional evidence and predicted effects for the risk

variant and its proxies ($R^2>0.8$) in the European population of the 1KGP3 project (1000 Genomes Project Consortium et al., 2015).

Four proxy variants were identified according to LDlink (Machiela and Chanock, 2015) (Supplementary Fig. S1). Notably, all variants influenced the *PIWIL4* gene, being rs508485 located in the 3'-UTR region, both the rs621425 and rs621922 polymorphisms in the 3' downstream region, and the rs504010 and rs1491388272 variants in intronic positions of the gene (Supplementary Table S5). Among these, rs504010 and rs508485 exhibited the strongest evidence of pathogenicity based on CADD, RegulomeDB, and FunSeq2 scoring (Supplementary Table S5). In addition, all variants overlapped with epigenetic marks such as H3K36me3 and H3K27me3, while rs621425 also overlapped with a CTCF-binding site (Supplementary Table S5), as indicated by ChIP-seq data from ENCODE (Davis et al., 2018). Further analyses predicted disruptions in transcription factor binding for the SNPs rs508485, rs504010, and rs621922, with affected transcription factors related to male infertility, spermatogenesis or having a predominant expression in the testis, including MEIS1 or GATA, among others (Supplementary Table S5). According to the transcriptome data of the GTEx project (GTEx Consortium, 2020), the proxies rs508485, rs621425, rs504010, and rs621922 were annotated as expression and splicing quantitative trait loci, thereby influencing the expression and splicing, respectively, of nearby genes in the testis (Supplementary Table S5, Supplementary Fig. S2). Furthermore, the Reactome pathway analysis linked rs508485, rs504010, and rs1491388272 to pathways such as 'gene silencing by RNA', 'piRNA biogenesis', and 'gene expression (transcription)' (Supplementary Table S5).

Finally, using the Open Targets Genetics Portal (Ghoussaini et al., 2021), we analysed which traits were associated with the rs508485 variant in the UK Biobank, FinnGen, and/or GWAS Catalog summary statistics repositories. Remarkably, the most associated trait with this polymorphism was the 'bioavailable testosterone levels', with a *P*-value of 1.40E-06 (Ruth et al., 2020) (Supplementary Fig. S3).

miRNA targets affected by the rs508485 polymorphism

Given the location of the rs508485 variant within the 3'-UTR region of *PIWIL4*, we initially used TargetScan, miRmap, and PolymiRTS to identify miRNAs with target sites that include this variant or are proximal to it. The identified miRNAs were hsa-miR-215-3p, hsa-miR-6773-3p, hsa-miR-4522-3p, hsa-miR-3664-5p, and hsa-miR-136-3p (Fig. 1). These computational tools assign a 'score' metric, that reflect the biological significance of each miRNA. Among these, the miRNAs hsa-miR-136-3p and hsa-miR-6773-3p exhibited the highest scores, indicative of their potential functional relevance. Subsequently, miRmap and RNAhybrid tools were employed to investigate the binding stability between the 3'-UTR sequence of the *PIWIL4* gene for each miRNA previously identified. All miRNAs exhibited increased stability in binding to the C allele of rs508485 compared to the T allele, except for hsa-miR-136-3p, which exhibited binding only in the presence of the T allele (Fig. 1). Furthermore, the difference in MFE values between the C and T alleles (ΔMFE) was computed. Remarkably low ΔMFE values were observed for all miRNAs, except for hsa-miR-136-3p, wherein, as previously mentioned, no binding occurred in the presence of the C allele, leading to *de novo* target formation in the presence of the T allele (Fig. 1).

Among all the miRNAs examined through the *in silico* analyses, rs508485 is located outside the recognition seed region in two of them (hsa-miR-3664-5p and hsa-miR4522-3p), thereby diminishing their relevance for this study. In the case of hsa-miR-6773-3p, while rs508485 localizes within the seed region, this miRNA lacks

Table 1. Associations of rs508485 genetic variant with severe SPGF subtypes when comparing with the control group under the considered models in the Iberian and German discovery cohorts and/or in the meta-analysis by the inverse variance method with the Iberian replication cohort.

SNP ID	Position (GRCh38)	EA	Pheno	Model	Iberian (discovery)			German (discovery)			Iberian (replication)			Meta-analysis		
					P	OR [95% CI]	OR [95% CI]	P	OR [95% CI]	P	OR [95% CI]	P	OR [95% CI]	P	OR [95% CI]	Q
rs508485	11:94621313	T	TESEneg	Recessive	3.01E-02	1.54 [1.04-2.28]	1.70 [1.13-2.54]	6.68E-01	1.17 [0.58-2.36]	1.09E-03	1.54 [1.19-2.00]	0.67	0.00			
			SCO	Additive	4.10E-02	1.35 [1.01-1.81]	1.36 [1.02-1.81]	4.36E-01	1.25 [0.72-2.17]	2.69E-03	1.34 [1.11-1.63]	0.96	0.00			

The results of the three independent studies are also shown. Significant P-values are highlighted in bold. CI, confidence interval; EA, effect allele; GRCh38, Genome Reference Consortium Human Build 38; HS, hypospermatogenesis; ID, identifier; MA, maturation arrest; NOA, non-obstructive azoospermia; OR, odds ratio; P, P-value; Pheno, phenotype; SCO, Sertoli cell-only; SPGF, spermatogenic failure; SNP, single-nucleotide polymorphism; TESEneg, negative outcome in testicular sperm extraction; sQTL, Splicing Quantitative Trait Locus.




Figure 1. Effect of the rs508485 polymorphism on the binding affinity of various microRNAs (miRNAs) to the 3'-untranslated region of the PIWIL4 gene. The target RNA and the different miRNAs sequences are shown, with binding base pairs in the seed region represented by vertical line. The polymorphism location within the target sequence is indicated by a black arrow. Predicted scores from TargetScan are displayed, where higher scores (less negative values) indicate stronger predicted repression. The figure also includes secondary structures, minimum free energy (MFE), and minimum free energy differences (Δ MFE) as calculated by the RNAhybrid tool. Lower MFE values indicate stronger miRNA binding affinity, while larger Δ MFE values reflect greater changes in miRNA binding affinity. Figure created in BioRender. Bossini-Castillo, L. (2025) <https://BioRender.com/118e878>.

established functionality and it is not expressed in the testis. However, both hsa-miR-215-3p and hsa-miR-136-3p include the polymorphism within their seed regions. Furthermore, they are expressed in the testis (Ludwig et al., 2016) and have been associated with the development of male infertility (Cheng et al., 2017; Zhang et al., 2021). Consequently, the functional assays were exclusively performed with hsa-miR-215-3p and hsa-miR-136-3p miRNAs, in conjunction with a scrambled miRNA as negative control; the sequences are shown in Supplementary Table S3.

Functional validation of the effect of the rs508485 polymorphism on miRNA binding affinity

To further investigate the binding interactions of both hsa-miR-215-3p and hsa-miR-136-3p miRNAs with the seed regions of the PIWIL4 3'-UTR according to presence of any of the two rs508485 alleles (T or C), we performed dual-luciferase reporter assays. Different miRNA concentrations (10, 25, and 50 nM) were tested to avoid non-specific effects resulting from cellular instability caused by excessively high miRNA concentrations.

The results showed that the hsa-miR-136-3p miRNA did not bind to the target sequence in the presence of the C allele, corroborating our previous *in silico* predictions (Fig. 2). Conversely, hsa-miR-215-3p exhibited a concentration-dependent decrease in luciferase activity, achieving ~60% specific inhibition at 50 nM compared to the scrambled miRNA control (Fig. 2). For the PIWIL4-3'-UTR region containing the T allele of the rs508485

polymorphism, has-miR-136-3p miRNA achieved ~75% specific inhibition at 50 nM, whereas hsa-miR-215-3p showed no significant impact on luciferase activity (Fig. 2). As anticipated, no variation in luciferase activity was observed for either miRNA when the mutated PIWIL4 3'-UTR sequence was used, due to the absence of binding sites resulting from the loss of the seed regions (Fig. 2).

Discussion

Understanding the genetic basis of SPGF is crucial for enhancing clinical management and genetic counselling for infertile patients (Sharma et al., 2021). SPGF often results from a complex interplay of genetic, epigenetic, and environmental factors (Cervan-Martin et al., 2020). While protein-coding genes were traditionally implicated in spermatogenesis, recent studies emphasize the importance of non-coding RNAs, particularly piRNAs and their associated PIWI proteins, in regulating male germ line development (Gou et al., 2017; Mann et al., 2023). Mutations and polymorphisms in both coding and non-coding genes can significantly impact spermatogenesis, highlighting the need for comprehensive genetic analyses to identify biomarkers for this multifaceted condition.

In the present study, we analysed the role of the rs508485 polymorphism, located in the 3'-UTR region of the PIWIL4 gene, in the predisposition to SPGF. This SNP has been previously




Figure 2. Influence of the rs508485 polymorphism on the binding affinity of hsa-miR-215-3p and hsa-miR-136-3p. (A) Schematic representation of psiCHECK-2 luciferase reporter constructs containing the 3'-untranslated regions (UTRs) of *PIWIL4*. The corresponding alleles are highlighted in red, microRNA (miRNA) seed sequences are in bold, and the mutated sequence is in brown. (B) Relative luciferase activity from dual-luciferase reporter assays. The percentage of luciferase activity normalized by the scrambled miRNA values versus the concentrations of hsa-miR-215-3p, hsa-miR-136-3p, and scrambled miRNAs (10, 25, and 50 nM) are represented for the presence of the T or C allele of rs508485 and a mutated sequence of *PIWIL4* 3'-UTR region. * $5E-02 \geq P > 5E-03$; ** $5E-03 \geq P > 5E-04$; *** $5E-04 \geq P$.

associated with certain patterns of male infertility (Gu *et al.*, 2010; Munoz *et al.*, 2014; Kamaliyan *et al.*, 2017, 2018). Our findings indicated that the T allele of rs508485 is a risk factor for SCO, the most severe form of male infertility, and may also predict unsuccessful TESE outcomes within a European genetic context. While the sample size of our discovery cohort provided sufficient statistical power to detect signals even in sub-phenotype groups of patients, the analysis performed in the replication cohort failed to replicate the observed associations due to a considerable reduction in the statistical power, particularly for the SPGF subgroups. Despite this, the allele effects remained consistent between the studied cohorts, with the T allele increasing the risk for SCO and TESE failure, suggesting an important role of the rs508485 genetic variant in the predisposition to the most severe forms of NOA. Indeed, the results of the meta-analysis showed an improvement in the statistical significance of the observed signals in both independent cohorts.

The results also shed light on the potential impact of the rs508485 genetic variant in SPGF development. The *in silico* analyses underscored its functional relevance, as indicated by its increased scores from multiple impact prediction algorithms and its correlation with active chromatin epigenetic marks. Additionally, the proximity of rs508485 to transcription factor binding sites, such as those for GATA (known to regulate Sertoli cell function and to maintain the spermatogonial stem cell niche; Kyronlahti *et al.*, 2011; Chen *et al.*, 2015) and MEIS1 (which is crucial for Sertoli cell-mediated regulation of male fertility; Sarkar *et al.*, 2021), further highlights its significance in spermatogenesis. Interestingly, rs508485 has been also reported to be associated with bioavailable testosterone levels (Ruth *et al.*, 2020). This steroid hormone, which is essential for testis development and normal spermatogenesis (Makela *et al.*, 2019), plays a critical role in male fertility and reproduction. Indeed, infertile patients with SPGF often exhibit reduced testosterone levels (Lardone *et al.*, 2018), and testosterone treatment is reported to induce elevated

piRNA expression in rats, suggesting that this hormone regulates testicular function through piRNA modulation (Kang et al., 2014). All of this functional evidence reinforces the association between rs508485 and specific patterns of male infertility observed in our study cohort.

Considering that rs508485 is located within the 3'-UTR region of *PIWIL4*, we hypothesized that this SNP affects the post-transcriptional regulation of *PIWIL4* by modifying miRNA binding affinity or accessibility. Dysregulation of gene expression by miRNAs, either through gene silencing by degrading mRNA or by disrupting their translation through base sequence complementarity with the target mRNA, has been linked to different diseases including neurodegenerative disorders, cancer, and cardiovascular conditions (Vishnoi and Rani, 2017). In the context of male infertility, miRNAs have been reported to play a critical role in the spermatogenic process (Procopio et al., 2017), and abnormal expression levels of miRNAs in semen samples have been associated with a reduced sperm count, low motility, and morphological abnormalities (Barbu et al., 2021). Our *in silico* analysis suggested that rs508485 influences miRNA binding affinity for *PIWIL4* transcripts. Specifically, hsa-miR-215-3p miRNAs form a more stable secondary structure when binding to the 3'-UTR region of *PIWIL4* in the presence of the C allele, whereas the T allele introduces a novel binding site for hsa-miR-136-3p. Dual-luciferase reporter assays confirmed these interactions, showing reduced luciferase activity associated with the C allele and hsa-miR-215-3p, and comparable reductions for the T allele and hsa-miR-136-3p.

Interestingly, hsa-miR-215-3p miRNA was found to be under-expressed in patients with SCO compared to fertile controls (Munoz et al., 2015; Zhang et al., 2021), and in individuals with Klinefelter syndrome (who are normally affected by NOA) (Sui et al., 2012). This miRNA is important for regulating cell cycle progression and cellular differentiation (Bueno and Malumbres, 2011; Jones et al., 2015; Vychytilova-Faltejskova and Slaby, 2019; Jia et al., 2022), both of which are critical events in


spermatogenesis. Notably, miR-215 is overexpressed in the testes during puberty and adulthood in boars and dears (Xu et al., 2018), suggesting that it may be crucial for spermatogonial differentiation in post-pubertal spermatogenesis.

In contrast, overexpression of hsa-miR-136-3p has been observed in SCO patients compared to unaffected men (Cheng et al., 2017). This miRNA is specifically expressed in spermatogonia (McIver et al., 2012) and, consequently, in the prepubertal testes of mammals (Liu et al., 2022). However, its precise role in male germ cell production remains unclear. It is important to note that *PIWIL4* is expressed from early stages of spermatogenesis (~14 weeks post-fertilization) until puberty (Guo et al., 2021). This timing aligns with the expression patterns of hsa-miR-215-3p and hsa-miR-136-3p miRNAs, as previously described (McIver et al., 2012; Xu et al., 2018; Guo et al., 2021; Liu et al., 2022) (Fig. 3). Taking all of the above into account, we strongly believe that slight alterations in the binding affinity of these miRNAs, caused by the risk allele of rs508485 in *PIWIL4*, lead to dysregulation of pathways involving this gene at crucial stages of germ cell development and differentiation. In this sense, we hypothesize that, in normal spermatogenesis, involving the rs508485 C allele, hsa-miR-136-3p does not recognize *PIWIL4* transcripts. However, hsa-miR-215-3p may downregulate *PIWIL4* expression at the onset of puberty, thereby promoting spermatogonial differentiation and supporting the progression of spermatogenesis (Fig. 3). On the contrary, the T allele introduces a novel binding site for hsa-miR-136-3p, potentially downregulating *PIWIL4* expression early germ cell stages in infants. Simultaneously, the T allele reduces hsa-miR-215-3p binding, impairing its regulatory role in repressing *PIWIL4* expression during puberty and, consequently, hindering the progression of spermatogenesis. This disruption may lead to the activation of transposable elements in pro-spermatogonia, driven by piRNA pathway dysfunction (Aravin et al., 2008; Kuramochi-Miyagawa et al., 2008), ultimately resulting in germ cell depletion and SCO (Fig. 3). Remarkably, the T allele effects over *PIWIL4* expression seem to be aligned with the miRNAs

C allele - normal germ cell development

- No interaction between hsa-miR-136 and the 3' UTR region of *PIWIL4*
- hsa-miR-215 downregulates *PIWIL4* expression, which may be crucial in spermatogonial differentiation

Germ cell development



T allele - SCO susceptibility

- *De novo* hsa-miR-136 seed
- Lower affinity of hsa-miR-215 to the 3' UTR region of *PIWIL4*
- *PIWIL4* deregulation may lead to germ cell death

Figure 3. Schematic representation of the hypothetical role of rs508485 in human male germ cell development. Cell types of the male germ cell lineage and the expression pattern of *PIWIL4*, hsa-miR-215-3p, and hsa-miR-136-3p are shown. PGC, primordial germ cell; SCO, Sertoli cell-only. Figure created in BioRender. Bossini-Castillo, L. (2025) <https://BioRender.com/thoyx6n>.

expression (up or down) effects in the SCO. This hypothesis is consistent with the association observed in our study cohort between this SNP and the SCO phenotype, and thus an unsuccessful TESE. Due to the complex aetiology of this disease, the rs508485 polymorphism cannot be considered a definitive biomarker for TESE success. However, its role in *PIWIL4* regulation via miRNA interactions highlights a broader genetic influence on spermatogenesis. From a clinical perspective, these findings may contribute to a more comprehensive patient assessment, where rs508485 genotyping could complement established clinical markers, improving the selection of NOA patients for sperm retrieval and enhancing pre-surgical counselling. In fact, disturbances in piRNAs expression profiles in testicular tissues of infertile men have been previously associated with TESE outcome (Cao *et al.*, 2018), and *MIWI2* knockout mice showed loss of germ cell lineages leading to SCO phenotype (Carmell *et al.*, 2007). Moreover, a recently published work also highlights that rare biallelic variants in protein-coding genes involved in piRNAs biogenesis led to defects in spermatogenesis and male infertility (Stallmeyer *et al.*, 2024). Thus, our study not only establishes rs508485 as a genetic susceptibility marker of SCO, but also identifies two miRNAs as potential novel regulators of *PIWIL4*, a critical player in retrotransposon control and spermatogenesis. Our findings align with previous reports suggesting an association between rs508485 and male infertility due to SPGF (Gu *et al.*, 2010; Munoz *et al.*, 2014; Kamaliyan *et al.*, 2017, 2018). However, our study provides additional evidence supporting a more specific role of this variant in SPGF, particularly in the SCO phenotype. While some studies have reported associations between rs508485 and SO (Gu *et al.*, 2010) or MA (Munoz *et al.*, 2014), our results did not support a broader implication of this SNP in other forms of SPGF. This divergence may be due to differences in cohort composition, genetic background, or phenotype characterization based on histopathological analysis across studies. Notably, our study includes the largest cohort of SPGF subtypes to date, which enhances the statistical power of our findings and reinforces their robustness compared to previous reports. Moreover, our functional assays strongly suggest that the impact of rs508485 is mediated through altered miRNA binding to *PIWIL4*, rather than affecting mRNA stability as previously hypothesized (Kamaliyan *et al.*, 2018). By providing functional validation of this mechanism and leveraging a more comprehensive cohort, our study clarifies the molecular role of this SNP and addresses the uncertainty in prior literature.

Nevertheless, this study has some limitations that should be considered when interpreting the results. First, approximately half of the NOA patients in our cohort lacked confirmatory histopathological diagnosis. In this regard, we would like to highlight that our key association was observed in the SCO subgroup, which represents the largest cohort of SCO patients evaluated so far. This strengthens the robustness of our findings despite the histopathological limitation. In any case, independent validation in well-characterized cohorts, including those from diverse ancestries, is warranted to confirm our findings. Additionally, while we provide functional evidence supporting the role of the rs508485 variant in *PIWIL4* regulation, our assays were conducted *in vitro* and focused on a single genetic polymorphism. Further studies using *in vivo* models and exploring additional genetic interactions are necessary to fully elucidate the molecular mechanisms underlying SPGF and to assess their potential implications for clinical practice.

In conclusion, our study provides valuable insights into the intricate molecular mechanisms underlying SCO, the most severe

form of SPGF. We establish a link between the rs508485 polymorphism and both SCO and TESE outcome, highlighting the significant role of common genetic variations in the pathogenesis of complex conditions such as idiopathic male infertility. Specifically, by functional validation, we demonstrate that the rs508485 genetic variant may influence miRNA binding to piRNA pathway genes like *PIWIL4*, shedding light on its potential contribution to spermatogenesis defects. These findings advance our understanding of SCO aetiology and suggest promising avenues for developing targeted therapies. By exploring the interaction between genetic predisposition and disease manifestation, our study contributes to the progress of personalized treatment strategies in male infertility. However, further research involving populations of different ancestries, larger cohorts and comprehensive functional analyses, is essential to unravel the biological mechanisms involved in the development of extreme patterns of male infertility and improve patient care outcomes.

Supplementary data

Supplementary data are available at *Human Reproduction* online.

Data availability

The data generated in this study are either contained in the article file and its [supplementary material](#) or can be made available upon reasonable request to the corresponding author.

Acknowledgements

We thank the National DNA Bank Carlos III (University of Salamanca, Spain) for supplying part of the control DNA samples from Spain and all the participants for their essential collaboration. The authors are also thankful for the support of the Unit of Excellence 'UNETE' from the University of Granada (reference UCE-PP2017-05). Also, we thank BioRender (through individual license) for providing some elements used in [Figs 1](#) and [3](#). This article is related to the PhD Doctoral Thesis of Sara González-Muñoz.

Authors' roles

R.J.P.-M. and F.D.C. were involved in the conception, design, and supervision of the study. S.G.-M., M.C.-M., A.G.-J., A.I.R.-M., and L.B.-C. participated in the methodology. S.G.-M., M.C.-M., A.G.-J., and A.I.R.-M. performed the formal analysis. S.G.-M., M.C.-M., A.G.-J., A.I.R.-M., and M.B. were involved in the interpretation of the data. N.G., J.A.C., M.C.G., A.C., M.M., M.A.V., A.E., V.M., M.L.G.-P., N.G.-G., E.S., C.G., F.Q.-F., S.G., D.A., L.M.-G., Y.O.-G., I.P.-C., G.S.P., A.A., I.S.P., O.L.-R., L.B., S.S., J.G., A.M.L., and S.L. were responsible for study subjects and data recruitment. S.G.-M. and M.C.-M. were involved in the original draft preparation. All authors critically revised and approved the final manuscript.

Funding

This work was supported by the Spanish Ministry of Economy and Competitiveness through the Spanish National Plan for Scientific and Technical Research and Innovation (grant: PID2020-120157RB-I00 funded by MICIU/AEI/10.13039/501100011033; and grant: PID2023-152215OB-I00 funded by MICIU/AEI/10.13039/501100011033 and ERDF, EU). M.C.-M. is beneficiary of a Juan de la Cierva 2022 postdoctoral fellowship (reference number JDC2022-

048561-I) funded by MCIN/AEI/10.13039/501100011033 and European Union NextGenerationEU/PRTR. S.L. received support from “Generalitat de Catalunya” (grant 2021SGR052). S.L. is sponsored by the “Researchers Consolidation Program” from the SNS-Dpt. Salut Generalitat de Catalunya (Exp. CES09/020). S.G.-M. was funded by grant ref. FPU23/02674. S.S. was supported by funds from the Portuguese Foundation for Science and Technology (FCT; 10.54499/DL57/2016/CP1363/CT0019).


Conflict of interest

None declared.


References

- 1000 Genomes Project Consortium; Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel JO, Marchini JL, McCarthy S, McVean GA, Abecasis GR et al. A global reference for human genetic variation. *Nature* 2015;**526**:68–74.
- Aravin AA, Hannon GJ, Brennecke J. The PIWI-piRNA pathway provides an adaptive defense in the transposon arms race. *Science* 2007a;**318**:761–764.
- Aravin AA, Sachidanandam R, Bourc’his D, Schaefer C, Pezic D, Toth KF, Bestor T, Hannon GJ. A piRNA pathway primed by individual transposons is linked to de novo DNA methylation in mice. *Mol Cell* 2008;**31**:785–799.
- Aravin AA, Sachidanandam R, Girard A, Fejes-Toth K, Hannon GJ. Developmentally regulated piRNA clusters implicate MILI in transposon control. *Science* 2007b;**316**:744–747.
- Barbu MG, Thompson DC, Suci N, Voinea SC, Cretoiu D, Predescu DV. The roles of microRNAs in male infertility. *Int J Mol Sci* 2021;**22**:2910.
- Bhattacharya A, Ziebarth JD, Cui Y. PolymiRTS Database 3.0: linking polymorphisms in microRNAs and their target sites with human diseases and biological pathways. *Nucleic Acids Res* 2014;**42**:D86–D91.
- Bueno MJ, Malumbres M. MicroRNAs and the cell cycle. *Biochim Biophys Acta* 2011;**1812**:592–601.
- Cao C, Wen Y, Wang X, Fang N, Yuan S, Huang X. Testicular piRNA profile comparison between successful and unsuccessful micro-TESE retrieval in NOA patients. *J Assist Reprod Genet* 2018;**35**:801–808.
- Carmell MA, Girard A, van de Kant HJ, Bourc’his D, Bestor TH, de Rooij DG, Hannon GJ. MIWI2 is essential for spermatogenesis and repression of transposons in the mouse male germline. *Dev Cell* 2007;**12**:503–514.
- Cervan-Martin M, Castilla JA, Palomino-Morales RJ, Carmona FD. Genetic landscape of nonobstructive azoospermia and new perspectives for the clinic. *J Clin Med* 2020;**9**:300.
- Cervan-Martin M, Gonzalez-Munoz S, Guzman-Jimenez A, Higuera-Serrano I, Castilla JA, Garrido N, Lujan S, Bassas L, Seixas S, Goncalves J et al. Changes in environmental exposures over decades may influence the genetic architecture of severe spermatogenic failure. *Hum Reprod* 2024;**39**:612–622.
- Cervan-Martin M, Tuttelmann F, Lopes AM, Bossini-Castillo L, Rivera-Egea R, Garrido N, Lujan S, Romeu G, Santos-Ribeiro S, Castilla JA et al. Immune and spermatogenesis-related loci are involved in the development of extreme patterns of male infertility. *Commun Biol* 2022;**5**:1220.
- Chen SR, Tang JX, Cheng JM, Li J, Jin C, Li XY, Deng SL, Zhang Y, Wang XX, Liu YX. Loss of Gata4 in Sertoli cells impairs the spermatogonial stem cell niche and causes germ cell exhaustion by attenuating chemokine signaling. *Oncotarget* 2015;**6**:37012–37027.
- Cheng YS, Chung CL, Chen CF, Lin YM. Differential expression of microRNAs and their messengerRNA targets in men with normal spermatogenesis versus Sertoli cell-only syndrome. *Urol Sci* 2017;**28**:42–49.
- Cocuzza M, Alvarenga C, Pagani R. The epidemiology and etiology of azoospermia. *Clinics (Sao Paulo)* 2013;**68**(Suppl 1):15–26.
- Davis CA, Hitz BC, Sloan CA, Chan ET, Davidson JM, Gabdank I, Hilton JA, Jain K, Baymuradov UK, Narayanan AK et al. The Encyclopedia of DNA elements (ENCODE): data portal update. *Nucleic Acids Res* 2018;**46**:D794–D801.
- Deng W, Lin H. MIWI, a murine homolog of PIWI, encodes a cytoplasmic protein essential for spermatogenesis. *Dev Cell* 2002;**2**:819–830.
- Dong S, Boyle AP. Predicting functional variants in enhancer and promoter elements using RegulomeDB. *Hum Mutat* 2019;**40**:1292–1298.
- Ghousaini M, Mountjoy E, Carmona M, Peat G, Schmidt EM, Hercules A, Fumis L, Miranda A, Carvalho-Silva D, Buniello A et al. Open Targets Genetics: systematic identification of trait-associated genes using large-scale genetics and functional genomics. *Nucleic Acids Res* 2021;**49**:D1311–D1320.
- Gou LT, Kang JY, Dai P, Wang X, Li F, Zhao S, Zhang M, Hua MM, Lu Y, Zhu Y et al. Ubiquitination-deficient mutations in human PIWI cause male infertility by impairing histone-to-protamine exchange during spermiogenesis. *Cell* 2017;**169**:1090–1104.e13.
- GTEX Consortium. The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science* 2020;**369**:1318–1330.
- Gu A, Ji G, Shi X, Long Y, Xia Y, Song L, Wang S, Wang X. Genetic variants in PIWI-interacting RNA pathway genes confer susceptibility to spermatogenic failure in a Chinese population. *Hum Reprod* 2010;**25**:2955–2961.
- Guo J, Sosa E, Chitiashvili T, Nie X, Rojas EJ, Oliver E, Plath K, Hotaling JM, Stukenborg J-B, Clark AT et al.; DonorConnect. Single-cell analysis of the developing human testis reveals somatic niche cell specification and fetal germline stem cell establishment. *Cell Stem Cell* 2021;**28**:764–778.e4.
- Hartig JV, Tomari Y, Forstemann K. piRNAs—the ancient hunters of genome invaders. *Genes Dev* 2007;**21**:1707–1713.
- He Z, Kokkinaki M, Pant D, Gallicano GI, Dym M. Small RNA molecules in the regulation of spermatogenesis. *Reproduction* 2009;**137**:901–911.
- Jia B, Zhang L, Ma F, Wang X, Li J, Diao N, Leng X, Shi K, Zeng F, Zong Y et al. Comparison of miRNA and mRNA expression in Sika deer testes with age. *Front Vet Sci* 2022;**9**:854503.
- Jones MF, Hara T, Francis P, Li XL, Bilke S, Zhu Y, Pineda M, Subramanian M, Bodmer WF, Lal A. The CDX1-microRNA-215 axis regulates colorectal cancer stem cell differentiation. *Proc Natl Acad Sci USA* 2015;**112**:E1550–E1558.
- Juliano C, Wang J, Lin H. Uniting germline and stem cells: the function of PIWI proteins and the piRNA pathway in diverse organisms. *Annu Rev Genet* 2011;**45**:447–469.
- Kamaliyan Z, Pouriamanesh S, Amin-Beidokhti M, Rezagholizadeh A, Mirfakhraie R. HIWI2 rs508485 polymorphism is associated with non-obstructive azoospermia in Iranian patients. *Rep Biochem Mol Biol* 2017;**5**:108–111.
- Kamaliyan Z, Pouriamanesh S, Soosanabadi M, Gholami M, Mirfakhraie R. Investigation of PIWI-interacting RNA pathway genes role in idiopathic non-obstructive azoospermia. *Sci Rep* 2018;**8**:142.
- Kang HJ, Moon MJ, Lee HY, Han SW. Testosterone alters testis function through regulation of piRNA expression in rats. *Mol Biol Rep* 2014;**41**:6729–6735.

- Klattenhoff C, Theurkauf W. Biogenesis and germline functions of piRNAs. *Development* 2008;**135**:3–9.
- Kozomara A, Birgaoanu M, Griffiths-Jones S. miRBase: from microRNA sequences to function. *Nucleic Acids Res* 2019;**47**:D155–D162.
- Kruger J, Rehmsmeier M. RNAhybrid: microRNA target prediction easy, fast and flexible. *Nucleic Acids Res* 2006;**34**:W451–W454.
- Kuramochi-Miyagawa S, Kimura T, Ijiri TW, Isobe T, Asada N, Fujita Y, Ikawa M, Iwai N, Okabe M, Deng W et al. Mili, a mammalian member of PIWI family gene, is essential for spermatogenesis. *Development* 2004;**131**:839–849.
- Kuramochi-Miyagawa S, Watanabe T, Gotoh K, Totoki Y, Toyoda A, Ikawa M, Asada N, Kojima K, Yamaguchi Y, Ijiri TW et al. DNA methylation of retrotransposon genes is regulated by PIWI family members MILI and MIWI2 in murine fetal testes. *Genes Dev* 2008;**22**:908–917.
- Kyronlahti A, Euler R, Bielinska M, Schoeller EL, Moley KH, Toppari J, Heikinheimo M, Wilson DB. GATA4 regulates Sertoli cell function and fertility in adult male mice. *Mol Cell Endocrinol* 2011;**333**:85–95.
- Lardone MC, Argandona F, Lorca M, Piottante A, Florez M, Palma C, Ebersperger M, Castro A. Leydig cell dysfunction is associated with post-transcriptional deregulation of CYP17A1 in men with Sertoli cell-only syndrome. *Mol Hum Reprod* 2018;**24**:203–210.
- Liu H, Khan IM, Liu Y, Khan NM, Ji K, Yin H, Wang W, Zhou X, Zhang Y. A comprehensive sequencing analysis of testis-born miRNAs in immature and mature indigenous Wandong cattle (*Bos taurus*). *Genes (Basel)* 2022;**13**:2185.
- Lorenz R, Bernhart SH, Honer Zu Siederdisen C, Tafer H, Flamm C, Stadler PF, Hofacker IL. ViennaRNA Package 2.0. *Algorithms Mol Biol* 2011;**6**:26.
- Ludwig N, Leidinger P, Becker K, Backes C, Fehlmann T, Pallasch C, Rheinheimer S, Meder B, Stahler C, Meese E et al. Distribution of miRNA expression across human tissues. *Nucleic Acids Res* 2016;**44**:3865–3877.
- Machiela MJ, Chanock SJ. LDlink: a web-based application for exploring population-specific haplotype structure and linking correlated alleles of possible functional variants. *Bioinformatics* 2015;**31**:3555–3557.
- Makela JA, Koskenniemi JJ, Virtanen HE, Toppari J. Testis development. *Endocr Rev* 2019;**40**:857–905.
- Mann JM, Wei C, Chen C. How genetic defects in piRNA trimming contribute to male infertility. *Andrology* 2023;**11**:911–917.
- Massart A, Lissens W, Tournaye H, Stouffs K. Genetic causes of spermatogenic failure. *Asian J Androl* 2012;**14**:40–48.
- McGeary SE, Lin KS, Shi CY, Pham TM, Bisaria N, Kelley GM, Bartel DP. The biochemical basis of microRNA targeting efficacy. *Science* 2019;**366**:eaav1741.
- McIver SC, Stanger SJ, Santarelli DM, Roman SD, Nixon B, McLaughlin EA. A unique combination of male germ cell miRNAs coordinates gonocyte differentiation. *PLoS One* 2012;**7**:e35553.
- Munoz X, Mata A, Bassas L, Larriba S. Altered miRNA signature of developing germ-cells in infertile patients relates to the severity of spermatogenic failure and persists in spermatozoa. *Sci Rep* 2015;**5**:17991.
- Munoz X, Navarro M, Mata A, Bassas L, Larriba S. Association of PIWIL4 genetic variants with germ cell maturation arrest in infertile Spanish men. *Asian J Androl* 2014;**16**:931–933.
- Oscanoa J, Sivapalan L, Gadaleta E, Dayem Ullah AZ, Lemoine NR, Chelala C. SNPnexus: a web server for functional annotation of human genome sequence variation (2020 update). *Nucleic Acids Res* 2020;**48**:W185–W192.
- Ozata DM, Gainetdinov I, Zoch A, O'Carroll D, Zamore PD. PIWI-interacting RNAs: small RNAs with big functions. *Nat Rev Genet* 2019;**20**:89–108.
- Procopio MS, de Avelar GF, Costa GMJ, Lacerda S, Resende RR, de Franca LR. MicroRNAs in Sertoli cells: implications for spermatogenesis and fertility. *Cell Tissue Res* 2017;**370**:335–346.
- Ruth KS, Day FR, Tyrrell J, Thompson DJ, Wood AR, Mahajan A, Beaumont RN, Wittmans L, Martin S, Busch AS et al.; Endometrial Cancer Association Consortium. Using human genetics to understand the disease impacts of testosterone in men and women. *Nat Med* 2020;**26**:252–258.
- Sarkar RK, Sen Sharma S, Mandal K, Wadhwa N, Kunj N, Gupta A, Pal R, Rai U, Majumdar SS. Homeobox transcription factor Meis1 is crucial to Sertoli cell mediated regulation of male fertility. *Andrology* 2021;**9**:689–699.
- Sharma A, Minhas S, Dhillon WS, Jayasena CN. Male infertility due to testicular disorders. *J Clin Endocrinol Metab* 2021;**106**:e442–e459.
- Stallmeyer B, Buhlmann C, Stakaitis R, Dicke AK, Ghieh F, Meier L, Zoch A, MacKenzie MacLeod D, Steingrover J, Okutman O et al. Inherited defects of piRNA biogenesis cause transposon derepression, impaired spermatogenesis, and human male infertility. *Nat Commun* 2024;**15**:6637.
- Sui W, Ou M, Chen J, Li H, Lin H, Zhang Y, Li W, Xue W, Tang D, Gong W et al. microRNA expression profile of peripheral blood mononuclear cells of Klinefelter syndrome. *Exp Ther Med* 2012;**4**:825–831.
- Tournaye H, Krausz C, Oates RD. Novel concepts in the aetiology of male reproductive impairment. *Lancet Diabetes Endocrinol* 2017;**5**:544–553.
- Vejnar CE, Zdobnov EM. MiRmap: comprehensive prediction of microRNA target repression strength. *Nucleic Acids Res* 2012;**40**:11673–11683.
- Vishnoi A, Rani S. MiRNA biogenesis and regulation of diseases: an overview. *Methods Mol Biol* 2017;**1509**:1–10.
- Vychytilova-Faltejskova P, Slaby O. MicroRNA-215: from biology to theranostic applications. *Mol Aspects Med* 2019;**70**:72–89.
- Ward LD, Kellis M. HaploReg v4: systematic mining of putative causal variants, cell types, regulators and target genes for human complex traits and disease. *Nucleic Acids Res* 2016;**44**:D877–D881.
- World Health Organization. *Infertility Is a Global Public Health Issue*. Geneva, Switzerland: World Health Organization, 2017.
- World Health Organization. *WHO Laboratory Manual for the Examination and Processing of Human Semen*, 6th edn. Geneva, Switzerland: World Health Organization, 2021.
- Xu XY, Wu D, Xu SY, Che LQ, Fang ZF, Feng B, Li J, Wu CM, Lin Y. Comparison of microRNA transcriptomes reveals differential regulation of microRNAs in different-aged boars. *Theriogenology* 2018;**119**:105–113.
- Zhang W, Zhang Y, Zhao M, Ding N, Yan L, Chen J, Gao L, Zhang G, Sun X, Gu Y et al. MicroRNA expression profiles in the seminal plasma of nonobstructive azoospermia patients with different histopathologic patterns. *Fertil Steril* 2021;**115**:1197–1211.



Supplementary Figure S1. Genomic context and putative functional proxy variants of the lead rs508485 polymorphism located in the 3'-untranslated region of PIWIL4 gene in the European population of the 1000 Genome Project Phase III. Each variant is labelled with its RegulomeDB score. The query variant (rs508485) is highlighted in blue, non-coding proxy variants are shown in yellow, and coding proxies are represented in red. Source: LDlink.



Supplementary Figure S2. Violin plot representations illustrating the quantitative trait locus (QTL) effects of *PIWIL4*-rs508485 in the testis. (A) Expression-QTL effects on RP11-867G2.6. (B) Splicing-QTL effects on RP11-867G2.5, RP11-867G2.6, and RP11-685N10.1. Source: GTEx Analysis Release V8.

Supplementary Table S1. Main clinical features of the 1516 infertile men included in the study.

Clinical feature	Discovery				Replication	
	Iberian (N = 627)		German (N = 647)		Iberian (N = 242)	
	N	Value	N	Value	N	Value
Age at diagnosis, years	230	36 (8)*	647	34 (7)*	235	39 (6)*
Non-obstructive azoospermia (NOA)	455	72.57%	325	50.23%	133	54.96%
Follicle stimulating hormone (FSH) levels, IU/L	227	15.20 (16.15)*	325	20.50 (15.40)*	102	13.65 (17.84)*
Luteinizing hormone (LH) levels, IU/L	184	5.60 (4.11)*	325	6.80 (4.90)*	86	5.87 (7.03)*
Testosterone levels, ng/ml	158	4.60 (2.22)*	324	13.85 (7.48)*	85	5.40 (3.34)*
Biopsy performed in NOA	257	56.48%	249	76.62%	70	52.63%
Sertoli cell-only (SCO)	104	40.47%	110	44.18%	28	40.00%
Meiotic arrest	49	19.07%	49	19.68%	12	17.14%
Hypospermatogenesis (HS)	43	16.73%	77	30.92%	3	4.29%
Negative outcome in testicular sperm extraction (TESEneg)	131	59.28%	127	57.47%	44	73.33%

Percentages refer to all individuals with available information for the variable.

* Median and interquartile range (IQR) are shown for those variables.

N, number of patients with available information; NOA, non-obstructive azoospermia; SCO, Sertoli cell-only; TESEneg, negative outcome in testicular sperm extraction.

Supplementary Table S2. Sequences of PIWIL4 3'-UTR region used in post-transcriptional gene silencing assays.

	3'-UTR sequence
T allele	ATAAGTGTGGCGTGATATTTGATGATAGATAAACAGAGTCTAATCCCACCCCA
C allele	ATAAGTGTGGCGTGATATTTGATGACAGATAAACAGAGTCTAATCCCACCCCA
Mutated	ATAAGTGTGGCGTGATATTT CAGCAGCTAG AAACAGAGTCTAATCCCACCCCA

T and C alleles of rs508485 polymorphism as well as 11 altered bases in mutated sequence are highlighted in bold.
 UTR, untranslated region.

Supplementary Table S3. Mimic microRNAs (miRNAs) used in functional assays and their respective sequences.

Mimic miRNA	Sequence
hsa-miR-215-3p	AUAACCGGAUUUCUUUACUGUCU
hsa-miR-136-3p	UCUGAGUAAACUCUGCUACUAC
Scrambled miRNA	AGAUGAGAAAGAUCUCCAACACU

Supplementary Table S4. Analysis of the allele frequencies of rs508485 genetic variant comparing subgroups of clinical phenotypes of male infertility against fertile controls under the additive, dominant, and recessive models.

SNP ID	Position (GRCh38)	EA	Pheno	Model	Iberian (discovery)			German (discovery)			Iberian (replication)			Meta-analysis		
					P	OR [95% CI]	I ²	P	OR [95% CI]	I ²	P	OR [95% CI]	I ²	P	OR [95% CI]	I ²
rs508485	11:94621313	T	SPGF	Additive	6.10E-01	1.04 [0.90-1.21]	2.66E-01	1.09 [0.94-1.25]	3.22E-01	0.89 [0.72-1.12]	5.32E-01	1.03 [0.94-1.13]	0.35	3.98		
				Dominant	8.30E-01	1.03 [0.80-1.31]	2.88E-01	1.14 [0.90-1.44]	3.54E-01	0.84 [0.59-1.21]	6.65E-01	1.03 [0.89-1.21]	0.40	0.00		
				Recessive	5.37E-01	1.08 [0.85-1.37]	4.55E-01	1.09 [0.87-1.38]	4.81E-01	0.87 [0.60-1.27]	5.52E-01	1.05 [0.90-1.22]	0.58	0.00		
		NOA		Additive	4.15E-01	1.07 [0.91-1.25]	8.91E-02	1.17 [0.98-1.41]	5.65E-01	0.92 [0.70-1.21]	1.75E-01	1.08 [0.97-1.20]	0.36	2.75		
				Dominant	9.00E-01	1.02 [0.78-1.33]	6.60E-02	1.33 [0.98-1.81]	1.69E-01	0.74 [0.48-1.14]	5.47E-01	1.06 [0.88-1.27]	0.09	59.36		
				Recessive	2.30E-01	1.17 [0.91-1.51]	3.57E-01	1.15 [0.86-1.54]	6.61E-01	1.11 [0.71-1.73]	1.18E-01	1.15 [0.96-1.38]	0.98	0.00		
		TESeneg		Additive	1.24E-01	1.23 [0.95-1.59]	8.18E-03	1.44 [1.10-1.88]	6.80E-01	0.91 [0.59-1.41]	1.11E-02	1.25 [1.05-1.48]	0.21	35.10		
				Dominant	7.60E-01	1.07 [0.69-1.67]	7.98E-02	1.53 [0.95-2.44]	2.73E-01	0.69 [0.35-1.35]	4.18E-01	1.13 [0.84-1.51]	0.16	46.11		
				Recessive	3.01E-02	1.54 [1.04-2.28]	1.03E-02	1.70 [1.13-2.54]	6.68E-01	1.17 [0.58-2.36]	1.09E-03	1.54 [1.19-2.00]	0.67	0.00		
		SCO		Additive	4.10E-02	1.35 [1.01-1.81]	3.82E-02	1.36 [1.02-1.81]	4.36E-01	1.25 [0.72-2.17]	2.69E-03	1.34 [1.11-1.63]	0.96	0.00		
				Dominant	1.09E-01	1.54 [0.91-2.62]	2.76E-02	1.81 [1.07-3.08]	4.09E-01	1.53 [0.56-4.23]	4.97E-03	1.66 [1.16-2.35]	0.90	0.00		
				Recessive	8.02E-02	1.47 [0.95-2.28]	2.53E-01	1.30 [0.83-2.04]	6.47E-01	1.22 [0.52-2.90]	3.69E-02	1.37 [1.02-1.83]	0.89	0.00		
		HS		Additive	7.37E-01	1.08 [0.70-1.66]	8.97E-01	0.98 [0.70-1.37]	NA	NA	9.19E-01	1.01 [0.78-1.32]	0.73	0.00		
				Dominant	3.10E-01	0.40 [0.69-3.28]	7.55E-01	1.09 [0.63-1.88]	NA	NA	4.04E-01	1.21 [0.77-1.89]	0.51	0.00		
				Recessive	6.46E-01	0.84 [0.40-1.76]	5.88E-01	0.85 [0.48-1.52]	NA	NA	4.78E-01	0.84 [0.54-1.34]	0.98	0.00		
		MA		Additive	4.40E-01	1.18 [0.78-1.79]	3.93E-01	1.20 [0.79-1.82]	2.21E-01	0.58 [0.24-1.40]	4.79E-01	1.11 [0.84-1.46]	0.31	13.97		
				Dominant	4.79E-01	1.30 [0.63-2.69]	6.00E-01	1.20 [0.60-2.41]	1.88E-01	0.42 [0.12-1.53]	7.41E-01	1.08 [0.68-1.73]	0.30	16.58		
				Recessive	5.73E-01	1.21 [0.63-2.34]	3.83E-01	1.33 [0.70-2.55]	4.77E-01	0.56 [0.11-2.77]	4.35E-01	1.19 [0.77-1.86]	0.61	0.00		

The results of the three independent studies and the meta-analysis are shown. Significant P-values are highlighted in bold. CI, confidence interval; EA, effect allele; GRCh38, Genome Reference Consortium Human Build 38; HS, hypospermatogenesis; ID, identifier; MA, maturation arrest; NA, non-available; NOA, non-obstructive azoospermia; OR, odds ratio; P, P-value; Pheno, phenotype; SCO, Sertoli cell-only; SNP, single-nucleotide polymorphism; SPGF, spermatogenic failure; TESeneg, negative outcome in testicular sperm extraction.

Supplementary Table S5. Comprehensive functional annotation of associated rs508485 variant and its proxies ($R^2 > 0.8$).

Proxies	Position (GRCh38)	R ²	Affected gene	Location	CADD	Functional prediction scores			Regulatory regions	Epigenetic marks	TFBS	eQTL (testis)	sQTL (testis)	Pathways
						Fun	Regulome	DB						
rs508485	11:94621313	1.00	PIWIL4	3'-Untranslated region	2.62	0.50	1f	-	H3K36me3	CCNT2; CDP; CEBPG; FOX; FOXD1; FOXL1; GATA; HDAC2; HOXA9; MAF; MEIS1; MRG1; PBX1	RP11-867G2.6	RP11-867G2.5; RP11-867G2.6; RP11-685N10.1	Gene silencing by RNA; piRNA biogenesis; Gene expression (transcription)	
rs621425	11:94622263	0.99	PIWIL4	3' Downstream	2.75	0.16	7	CTCF binding site	H3K36me3; H3K27me3; H3K27ac	NF1	RP11-867G2.6	RP11-867G2.5; RP11-867G2.6; RP11-685N10.1	-	
rs504010	11:94620853	0.99	PIWIL4	Intronic	8.42	0.34	1f	-	H3K36me3; H3K27me3; H3K4me1	AIRE; CART1; CDP; CUX2; DBX2; EMX2; EVI1; GATA; HNF1A; HNF1B; HOXB7; HOXC6; HOXD8; MSX1; NCX; NKX6; PAX4; POU2F2; PRRX2; SOX; VAX2	RP11-867G2.6; RP11-685N10.1	RP11-867G2.5; RP11-867G2.6; RP11-685N10.1	Gene silencing by RNA; piRNA biogenesis; Gene expression (transcription)	
rs621922	11:94622148	0.98	PIWIL4	3' Downstream	0.90	0.00	6	-	H3K36me3; H2BK12ac; H2AK5ac; H3K27me3	AP1; ETS; GATA; HDAC2; HNF1; MEF2; P300; PAX5; PBX1; SPI1; SPIB; STAT; TFII	RP11-867G2.6; MRE11	RP11-867G2.5; RP11-867G2.6	-	
rs1491388272	11:94610226	0.91	PIWIL4	Intronic	-	-	7	-	H3K36me3; H3K9me3	-	-	-	Gene silencing by RNA; piRNA biogenesis; Gene expression (transcription)	

Transcription factor binding sites (TFBS) related to male infertility, spermatogenesis, or having a predominant expression in the testis are marked in bold. eQTL, Expression Quantitative Trait Locus; GRCh38, Genome Reference Consortium Human Build 38; sQTL, Splicing Quantitative Trait Locus; piRNA, PIWI-interacting RNAs.