

Predominance of KPC-3 in a Survey for Carbapenemase-Producing *Enterobacteriaceae* in Portugal

Vera Manageiro,^{a,b} Eugénia Ferreira,^a Joana Almeida,^a Stephanie Barbosa,^a Constança Simões,^a Antibiotic Resistance Surveillance Program in Portugal (ARSIP), Robert A. Bonomo,^c Manuela Caniça^a

National Reference Laboratory of Antibiotic Resistances and Healthcare Associated Infections (NRL-AMR-HAI), Department of Infectious Diseases, National Institute of Health Dr. Ricardo Jorge, Lisbon, Portugal^a; CECA-ICETA, Centro de Estudos de Ciência Animal, Universidade do Porto, Porto, Portugal^b; Research Service, Louis Stokes Cleveland VA Medical Center, Department of Medicine, Pharmacology, Molecular Biology, and Microbiology, Case Western Reserve University, Cleveland, Ohio, USA^c

Among the 2,105 *Enterobacteriaceae* tested in a survey done in Portugal, 165 were nonsusceptible to carbapenems, from which 35 (26 *Klebsiella pneumoniae*, 3 *Escherichia coli*, 2 *Enterobacter aerogenes*, and 3 *Enterobacter cloacae* isolates and 1 *Klebsiella oxytoca* isolate) were confirmed to be carbapenemase producers by the presence of 30 *Tn4401d-bla*_{KPC-3}, 4 *intI3-bla*_{GES-5}, and one *intI1-bla*_{VIM-2} gene, alone or in combination with other *bla* genes. The dissemination of *bla*_{KPC-3} gene carried by an IncF plasmid suggests lateral gene transfer as a major mechanism of dissemination.

Carbapenem-nonsusceptible *Enterobacteriaceae* are increasingly reported worldwide, mainly because of the acquisition of carbapenemase-encoding genes located on highly mobile genetic elements (e.g., plasmids and/or integrons) that facilitate their horizontal spread (1, 2). In Portugal, little is known regarding the molecular epidemiology of carbapenemase-producing *Enterobacteriaceae* (CPE). Indeed, only sporadic isolates or single hospital cases have been described (3), namely, a VIM-2-producing *Klebsiella oxytoca* isolate (4), 2 VIM-34-producing *Klebsiella pneumoniae* isolates (5), and 6 *K. pneumoniae* carbapenemase (KPC)-3-producing *Enterobacteriaceae* isolates (6). The objectives of this study were to (i) characterize a large number of CPE isolates recovered from different Portuguese health care institutions, (ii) determine the potential diversity of these isolates, and (iii) describe the respective plasmids harboring carbapenemase-encoding genes in an international context.

This study included 2,105 consecutive and nonrepetitive clinical *Enterobacteriaceae* isolates that were collected between April 2006 and February 2013 and sent to the National Reference Laboratory of Antimicrobial Resistances at the National Institute of Health, as an integrative part of the Antimicrobial Resistance Surveillance Program in Portugal (ARSIP) laboratory. ARSIP is a voluntary Portuguese surveillance program that continually monitors the susceptibilities of pathogens of clinical importance against several antibiotics. Thirteen Portuguese hospitals located in five different regions (North, Central, Lisbon and Tagus Valley, South, and Autonomous Region of Madeira) participated in this survey.

The screening of carbapenem susceptibility, performed by the disk diffusion method, according to the French Society for Microbiology (<http://www.sfm-microbiologie.org/>), revealed that 165 (7.8%) isolates were ertapenem-nonsusceptible, and these were selected for further analysis. In addition, clinical isolates showing an inhibition of carbapenemase, as evidenced by the synergy between carbapenems (imipenem, meropenem, and/or ertapenem) and 3-aminophenylboronic acid or EDTA, were considered presumptive carbapenemase producers from class A or B, respectively (7).

Thirty-five isolates suspected to produce CPE (35/168 [20.8%]) demonstrated a positive synergy between carbapenems and 3-aminophenylboronic acid ($n = 34$) or with EDTA

($n = 1$), indicative of serine or metallo- β -lactamase production, respectively. The majority of those isolates were collected from urine samples (54.3%) from elderly (≥ 65 years old) male patients (54.3%), admitted to the emergency room/ambulatory (22.9%), internal medicine (17.1%), or surgery (17.1%) wards. The diminished susceptibilities to carbapenems among the remaining 130 ertapenem-nonsusceptible isolates (59 *Klebsiella* spp., 28 *Enterobacter* spp., 23 *Escherichia coli*, 11 *Morganella morganii*, 8 *Proteus mirabilis*, and 1 *Serratia marcescens*) suggested a decrease in the outer membrane permeability associated with the weak hydrolytic activity of carbapenems due to the expression of a β -lactamase, such as extended-spectrum β -lactamases (ESBLs), and/or AmpC overexpression (7).

The MICs of the 35/168 isolates were obtained by the reference microdilution broth method (8, 9), according to the EUCAST guidelines (http://www.eucast.org/clinical_breakpoints/). Tigecycline and ciprofloxacin were potent against 57.1% and 28.6% of the isolates, respectively, but colistin was the only antibiotic effective against all CPE isolates; all isolates were multidrug resistant (MDR), i.e., presented reduced susceptibility to three or more structurally unrelated antibiotics (10) (Table 1).

PCR amplification and DNA sequencing were applied to the 165 ertapenem-nonsusceptible isolates to identify the presence or confirm the absence of the carbapenemase-encoding genes *bla*_{KPC} and *bla*_{GES} (class A), *bla*_{IMP}, *bla*_{VIM}, and *bla*_{NDM} (class B), and *bla*_{OXA-48} (class D) (11). Among the 165 isolates, only the 35 identified as carbapenemase-producing isolates were confirmed to be

Received 16 December 2014 Returned for modification 11 January 2015
Accepted 13 March 2015

Accepted manuscript posted online 16 March 2015

Citation Manageiro V, Ferreira E, Almeida J, Barbosa S, Simões C, Antibiotic Resistance Surveillance Program in Portugal (ARSIP), Bonomo RA, Caniça M. 2015. Predominance of KPC-3 in a survey for carbapenemase-producing *Enterobacteriaceae* in Portugal. *Antimicrob Agents Chemother* 59:3588–3592. doi:10.1128/AAC.05065-14.

Address correspondence to Manuela Caniça, manuela.canica@insa.min-saude.pt.

Copyright © 2015, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AAC.05065-14

TABLE 1 MIC data and susceptibilities of 35 CPE isolates

Antibiotic ^a	All isolates (n = 35)										KPC-3-producing isolates (n = 30)										GES-5-producing isolates (n = 4)										VIM-2-producing isolates (n = 1)	
	MIC ₅₀		MIC ₉₀		MIC range		%R		%IR		MIC ₅₀		MIC ₉₀		MIC range		%R		%IR		MIC (μg/ml) for strain:		MIC for Ko9864		Susceptibility ^d							
																					Kp15587	Kp15743	Kp17060	Kp17061	%R	%IR	(μg/ml)	Susceptibility ^d				
AMX	>256	>256	>256	>256	>256	100.0	100.0	>256	>256	>256	100.0	100.0	>256	>256	100.0	100.0	>256	>256	>256	>256	>256	>256	>256	100.0	100.0	>256	R					
AMC	>256	>256	>256	>256	>256	100.0	100.0	>256	>256	>256	100.0	100.0	>256	>256	100.0	100.0	>256	>256	>256	>256	>256	>256	>256	100.0	100.0	>256	R					
TIC	>256	>256	>256	>256	>256	100.0	100.0	>256	>256	>256	100.0	100.0	>256	>256	100.0	100.0	>256	>256	>256	>256	>256	>256	>256	100.0	100.0	>256	R					
CXM	>256	>256	64 to >256	64 to >256	64 to >256	100.0	100.0	>256	>256	64 to >256	100.0	100.0	>256	>256	100.0	100.0	>256	>256	>256	>256	>256	>256	>256	100.0	100.0	>256	R					
FOX ^c	128	>128	16 to >128	16 to >128	16 to >128	100.0	100.0	128	>128	16 to 128	100.0	100.0	>128	128	128	>128	100.0	100.0	>128	>128	>128	>128	>128	100.0	100.0	>128	NW ^d					
CAZ	128	512	64 to 512	64 to 512	64 to 512	100.0	100.0	128	512	128 to >512	100.0	100.0	256	256	128	128	256	128	256	128	256	128	256	100.0	100.0	128	R					
CAZ-CLA	128	256	16 to 512	16 to 512	16 to 512	100.0	100.0	256	256	16 to 512	100.0	100.0	32	32	16	16	32	32	16	32	32	16	32	100.0	100.0	256	R					
CTX	256	>256	64 to >512	64 to >512	64 to >512	100.0	100.0	256	>256	64 to >256	100.0	100.0	64	64	64	64	64	64	64	64	64	64	64	100.0	100.0	256	R					
CTX-CLA	64	256	8 to >256	8 to >256	8 to >256	100.0	100.0	64	256	8 to >256	100.0	100.0	64	64	32	32	64	64	32	64	64	64	64	100.0	100.0	32	R					
ATM	256	512	32 to >512	32 to >512	32 to >512	100.0	100.0	256	512	64 to 512	100.0	100.0	256	256	256	256	128	32	32	32	32	32	32	100.0	100.0	128	R					
FEP	32	>256	8 to >256	8 to >256	8 to >256	100.0	100.0	32	>256	16 to >256	100.0	100.0	8	16	16	8	8	8	8	8	8	8	8	100.0	100.0	8	R					
IPM	8	64	0.5 to >512	0.5 to >512	0.5 to >512	48.6	85.7	8	32	1 to 128	46.7	90.0	>512	512	1	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	50.0	50.0	16	R					
MEM	32	64	0.25 to >128	0.25 to >128	0.25 to >128	85.7	97.1	32	64	0.25 to >128	90.0	96.7	128	128	8	4	4	4	4	4	4	4	4	50.0	100.0	64	R					
DOR	8	32	0.5 to 64	0.5 to 64	0.5 to 64	82.9	91.4	8	16	1 to 64	86.7	96.7	64	64	1	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	50.0	50.0	8	R					
ERT	32	64	1 to >64	1 to >64	1 to >64	91.4	100.0	32	64	1 to >64	90.0	100.0	32	>64	16	8	8	8	8	8	8	8	8	100.0	100.0	64	R					
CIP	32	>32	≤0.06 to >32	≤0.06 to >32	≤0.06 to >32	62.9	71.4	4	16	≤0.06 to >32	56.7	66.7	>32	>32	32	>32	>32	>32	>32	>32	>32	>32	>32	100.0	100.0	32	R					
GEN	32	512	4 to 512	4 to 512	4 to 512	97.1	100.0	32	256	4 to 256	96.7	100.0	512	512	512	512	512	512	512	512	512	512	512	100.0	100.0	256	R					
TMP	>1,024	>1,024	>1,024	>1,024	>1,024	100.0	100.0	>1,024	>1,024	>1,024	100.0	100.0	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	100.0	100.0	>1,024	R				
CST	0.25	1	≤0.03 to 2	≤0.03 to 2	≤0.03 to 2	0.0	0.0	0.25	1	≤0.03 to 2	0.0	0.0	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.0	0.0	0.5	S					
TGC	1	4	≤0.03 to 8	≤0.03 to 8	≤0.03 to 8	20.0	42.9	1	4	≤0.03 to 8	23.3	33.3	2	2	2	2	2	2	2	2	2	2	2	0.0	100.0	2	I					

^a AMX, amoxicillin; AMC, amoxicillin-clavulanic acid; TIC, ticarcillin; CXM, cefuroxime; FOX, cefoxitin; CAZ, ceftazidime; CAZ-CLA, ceftazidime-clavulanic acid; CTX, cefotaxime; CTX-CLA, ceftaxime-clavulanic acid; ATM, aztreonam; FEP, ceftipime; IPM, imipenem; MEM, meropenem; DOR, doripenem; ERT, ertrapenem; CIP, ciprofloxacin; GEN, gentamicin; TMP, trimethoprim; CST, colistin; and TGC, tigecycline.

^b R, resistant; IR, nonsusceptible.

^c Epidemiological breakpoint, according to EUCAST (http://www.eucast.org/mic_distributions_scoffs/).

^d NW^d, non-wild type; S, susceptible; I, intermediate.

TABLE 2 Characteristics of 35 CPE isolates detected in this study and of the 10 transconjugant strains obtained^a

Species (no.) encountered by carbapenemase type ^b	MLST (no.)	Hospital code/yr of isolation (no.)	Other β -lactamase(s) detected ^c
KPC-3 (<i>Tn4401b</i>)			
<i>E. aerogenes</i> (3)		<u>A/2012</u> , E/2011, H/2013	TEM-1, (OXA-30), (CTX-M-15)
<i>E. cloacae</i> (2)		E/2010, E/2012	ACT-type, (TEM-1), (OXA-30)
<i>E. coli</i> (<i>n</i> = 3)	ST58	G/2013	TEM-1
<i>K. pneumoniae</i> (22)	ST11 (2)	E/2012, E/2013	TEM-1, SHV-11, (OXA-30), (CTX-M-15)
	ST14 (3)	<u>F/2010</u> , G/2012, G/2013	TEM-1, SHV-1
	ST15 (4)	<u>G/2012</u> (2), G/2012 (1), G/2013 (1)	TEM-1, SHV-1, CTX-M-15
	ST34 (1)	<u>E/2011</u>	TEM-1, SHV-26
	ST59 (1)	E/2011	TEM-1
	ST147 (4)	<u>E/2012</u> (2), E/2012 (2)	TEM-1, SHV-11
	ST416 (1)	B/2011	TEM-1, SHV-14, OXA-30, CTX-M-15
	ST698 (1)	<u>G/2012</u>	TEM-1, SHV-11, CTX-M-15
	ST960 (1) ^d	<u>I/2011</u>	TEM-1, SHV-164
	ST1138 (4) ^d	<u>D/2011</u> (1), D/2011 (1), E/2011 (2)	TEM-1, SHV-36
GES-5 (<i>intI3</i>)			
<i>K. pneumoniae</i> (4)	ST231	C/2012 (2), C/2013 (2)	TEM-1, SHV-12, SHV-1
VIM-2 (<i>intI1</i>)			
<i>K. oxytoca</i> (1)		H/2010	SHV-12

^a Isolates that successfully transferred the KPC-3 to recipient strains are underlined. All clinical isolates and their respective transconjugant isolates contained KPC-3- and TEM-1-harboring IncF(FII) plasmids.

^b Parentheses indicate the mobile genetic element associated with the respective carbapenemase-encoding gene.

^c Parentheses indicate the variable presence of an antibiotic resistance determinant.

^d MLST first described here.

CPE by the presence of 30 *bla*_{KPC-3} and 4 *bla*_{GES-5} genes and 1 *bla*_{VIM-2} gene, alone or in combination with other *bla* genes (Table 2) (11).

An analysis of the genetic elements associated with the CPE-encoding genes showed that all the *bla*_{KPC-3} genes were flanked by IS*Kpn7* on the right and IS*Kpn6* on the left (12) and a Tn3-like Tn4401 transposon, with a 68-bp deletion upstream of *bla*_{KPC-3}, corresponding to the isoform d. This uncommon isoform has been described so far only in KPC-producing *K. pneumoniae* strains with high-level carbapenem resistance in the United States and Canada (13–15).

Regarding the *bla*_{GES-5} genes, the PCR results and sequence analysis, using previously described primers (11), revealed the amplification of the *intI3-bla*_{GES-5} region, where the *intI3* gene was located adjacent to the *bla*_{GES-5} gene in a configuration previously reported for the *bla*_{GES-1} gene (16, 17). *intI3* was targeted and amplified with a set of primer pairs (5'-GCAAGTGGGTGGCGA ATG-3' and 5'-CTGAAGTCGAGGGTTTTCTG-3') designed in this study (with initial denaturation at 94°C for 5 min, 30 cycles of denaturation at 94°C for 30 s, annealing at 58°C for 60 s, and extension at 72°C for 60 s, and a final extension at 72°C for 5 min). This study represents one of the very few reports of the acquisition of carbapenemase-encoding genes mediated by a class 3 integron (18, 19), and to our knowledge, those are the first such genes to have been found in CPE isolates. Notable is the fact that two isolates (Kp17060 and Kp17061, Table 1) were collected on the same day from patients admitted in the same hospital room, suggesting that organisms were transferred through direct contact. Also, we confirmed the location of the *bla*_{VIM-2} gene in a class 1 integron, as previously described (4).

In our study, among the 15 isolates with CPE-encoding genes randomly selected to be tested for transferability by broth

mating-out assays using recipient rifampin-resistant (Rif^r) or streptomycin-resistant (Str^r) *E. coli* C600 and sodium azide-resistant (NaN₃^r) *E. coli* J53 strains (11), we obtained 10 (66.7%) transconjugants with *bla*_{KPC-3}-harboring plasmids. Using PCR-based replicon typing to type the resistance plasmids (20), we observed the transfer of an IncF group harboring the *bla*_{KPC-3} and *bla*_{TEM-1} genes within the recipient NaN₃^r *E. coli* J53 (*n* = 9) and Rif^r *E. coli* C600 (*n* = 1) strains. The location of the *bla*_{KPC} gene has been observed in plasmids of various sizes, belonging to the IncF, IncL/M, and IncN plasmid types, and also on small rolling circle-replicating (RCR) plasmids that are not self-transmissible but can mobilized in *trans* by coresident conjugative plasmids (2). In this study, we identified the IncF plasmid in different species, such as *Klebsiella* spp. and *Enterobacter* spp., suggesting that the dissemination of the *bla*_{KPC-3} gene is due to lateral gene transfer rather than clonal spread. In general, the MICs of the transconjugants showed similar susceptibility profiles as those of the donor strains with nonsusceptibility to meropenem, ertapenem, cefotaxime, ceftazidime, gentamicin, and trimethoprim, as well as those with susceptibility to colistin and tigecycline. Regarding imipenem (*n* = 7), doripenem (*n* = 9), and ciprofloxacin (*n* = 9), the majority of the transconjugants were susceptible in comparison to the clinical isolates, with MICs from 8 mg/liter to 1 mg/liter for imipenem, 2 mg/liter to 0.25 mg/liter for doripenem, and 0.125 mg/liter to \leq 0.06 mg/liter for ciprofloxacin.

Multilocus sequence typing of *K. pneumoniae* (*n* = 26) isolates (<http://bigsdw.web.pasteur.fr/klebsiella/klebsiella.html>) showed a high degree of genetic diversity, as they were from distinct sequence types (STs), namely, ST14, ST15, ST34, ST59, ST147, ST416, and ST698, and from the two novel STs ST960 and ST1138 (Table 2). Overall, 10 STs were observed to be associated with KPC production, with ST147 (*n* = 4), ST1138 (*n* = 4), and ST15 (*n* =

4) being the most prevalent types. ST147 is an important clone associated with this resistance mechanism worldwide, namely due to VIM-, KPC-, and NDM-producing *K. pneumoniae* isolates (21, 22), while ST1138 was first described here in 4 KPC-3-producing *K. pneumoniae* isolates coexpressing TEM-1 and SHV-36 (Table 2). Interestingly, isolates belonging to the well-known lineage of *K. pneumoniae* ST258, which plays a major role in the global spread of KPC carbapenemases (23), were not found. Regarding KPC-producing *E. coli* ($n = 3$) (<http://mlst.warwick.ac.uk/mlst/dbs/Ecoli>), all isolates belonged to ST58.

In conclusion, this study provides new data regarding the molecular epidemiology of CPE in Portugal, with the emergence of the carbapenemase KPC-3. Indeed, this study suggests that the dissemination of the *bla*_{KPC-3} gene occurs due to the dispersion of Tn4401d, carried by an IncF plasmid and spread both by polyclonal *K. pneumoniae* and by *Enterobacter aerogenes*, *Enterobacter cloacae*, and *E. coli* clinical strains.

ACKNOWLEDGMENTS

V. Manageiro was supported by grant SFRH/BPD/77486/2011 from the Fundação para a Ciência e a Tecnologia, Lisbon, Portugal. J. Almeida was supported by grant BRJ-SUB/01/2012 from the National Institute of Health Dr. Ricardo Jorge, Lisbon, Portugal. R. A. Bonomo was supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under awards R01 AI100560 and R01 AI063517, and by the Cleveland Department of Veterans Affairs, the Veterans Affairs Merit Review Program, and the Geriatric Research Education and Clinical Center VISN 10.

We thank the Fundação para a Ciência e a Tecnologia (FCT) for project grant PEst-OE/AGR/UI0211/2011-2014, Strategic Project U1211-2011-2014. We also thank the laboratories participating in the Antibiotic Resistance Surveillance Program in Portugal (ARSIP) for sending isolates to the NRL-AMR/HAI at the NIH: Centro Hospitalar de Vila Real/Peso da Régua-EPE (A. P. Castro), Centro Hospitalar de Vila Nova de Gaia/Espinho-EPE (P. Lopes), Centro Hospitalar Póvoa de Varzim e Vila do Conde-EPE (F. Fonseca), Unidade Local de Saúde do Alto Minho (S. Vieira), and IPO-Porto (M. A. Guimarães) in the North region; Hospital São Teotónio, Viseu (J. Ribeiro) and Centro Hospitalar de Coimbra (H. Oliveira) in the Central region; Centro Hospitalar Lisboa Central (M. Pinto), Hospital Garcia de Orta-EPE (J. Diogo), Centro Hospitalar Barreiro-Montijo-EPE (A. Jesus), Hospital Prof. Doutor Fernando Fonseca-EPE (L. Sancho), and Hospital Vila Franca de Xira (M. Rodrigues) in the Lisbon and Tagus Valley region; Centro Hospitalar Barlavento Algarvio-EPE (T. Vaz) in the South region; and Centro Hospitalar Funchal (T. Afonso) in the Autonomous Region of Madeira.

REFERENCES

- Nordmann P, Naas T, Poirel L. 2011. Global spread of carbapenemase-producing *Enterobacteriaceae*. *Emerg Infect Dis* 17:1791–1798. <http://dx.doi.org/10.3201/eid1710.110655>.
- Carattoli A. 2013. Plasmids and the spread of resistance. *Int J Med Microbiol* 303:298–304. <http://dx.doi.org/10.1016/j.ijmm.2013.02.001>.
- Glasner C, Albiger B, Buist G, Tambić Andrasević A, Canton R, Carmeli Y, Friedrich AW, Giske CG, Glupczynski Y, Gniadkowski M, Livermore DM, Nordmann P, Poirel L, Rossolini GM, Seifert H, Vatopoulos A, Walsh T, Woodford N, Donker T, Monnet DL, Grundmann H, the European Survey on Carbapenemase-Producing *Enterobacteriaceae* (EuSCAPE) Working Group. 2013. Carbapenemase-producing *Enterobacteriaceae* in Europe: a survey among national experts from 39 countries. *Euro Surveill* 18:pil=20525. <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=20525>.
- Conceição T, Brizio A, Duarte A, Barros R. 2005. First isolation of *bla*_{VIM-2} in *Klebsiella oxytoca* clinical isolates from Portugal. *Antimicrob Agents Chemother* 49:476. <http://dx.doi.org/10.1128/AAC.49.1.476.2005>.
- Rodrigues C, Novais A, Machado E, Peixe L. 2014. Detection of VIM-34, a novel VIM-1 variant identified in the intercontinental ST15 *Klebsiella pneumoniae* clone. *J Antimicrob Chemother* 69:274–275. <http://dx.doi.org/10.1093/jac/dkt314>.
- Manageiro V, Ferreira E, Louro D, The Antimicrobial Resistance Surveillance Program in Portugal (ARSIP), Caniça M. 2013. Polyclonal KPC-3-producing *Enterobacteriaceae* in Portugal, p 212–216. In Mendez-Vilas A (ed), *Worldwide research efforts in the fighting against microbial pathogens: from basic research to technological developments*. Brown-Walker Press, Boca Raton, FL.
- Giske CG, Gezelius L, Samuelsen Ø, Warner M, Sundsfjord A, Woodford N. 2011. A sensitive and specific phenotypic assay for detection of metallo-β-lactamases and KPC in *Klebsiella pneumoniae* with the use of meropenem disks supplemented with aminophenylboronic acid, dipicolinic acid and cloxacillin. *Clin Microbiol Infect* 17:552–556. <http://dx.doi.org/10.1111/j.1469-0691.2010.03294.x>.
- European Committee on Antimicrobial Susceptibility Testing (EUCAST) Steering Committee. 2006. EUCAST technical note on tigecycline. *Clin Microbiol Infect* 12:1147–1149. <http://dx.doi.org/10.1111/j.1469-0691.2006.01578.x>.
- European Committee for Antimicrobial Susceptibility Testing (EUCAST) of the European Society of Clinical Microbiology and Infectious Diseases (ESCMID). 2003. Determination of minimum inhibitory concentrations (MICs) of antibacterial agents by broth dilution. *Clin Microbiol Infect* 9:ix–xv. <http://dx.doi.org/10.1046/j.1469-0691.2003.00790.x>.
- Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, Harbarth S, Hindler JF, Kahlmeter G, Olsson-Liljequist B, Paterson DL, Rice LB, Stelling J, Struelens MJ, Vatopoulos A, Weber JT, Monnet DL. 2011. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 18:268–281. <http://dx.doi.org/10.1111/j.1469-0691.2011.03570.x>.
- Manageiro V, Ferreira E, Caniça M, Manaia CM. 2014. GES-5 among the β-lactamases detected in ubiquitous bacteria isolated from aquatic environment samples. *FEMS Microbiol Lett* 351:64–69. <http://dx.doi.org/10.1111/1574-6968.12340>.
- Giani T, D'Andrea MM, Pecile P, Borgianni L, Nicoletti P, Tonelli F, Bartoloni A, Rossolini GM. 2009. Emergence in Italy of *Klebsiella pneumoniae* sequence type 258 producing KPC-3 carbapenemase. *J Clin Microbiol* 47:3793–3794. <http://dx.doi.org/10.1128/JCM.01773-09>.
- Mataseje LF, Boyd DA, Willey BM, Prayitno N, Kreiswirth N, Gelosia A, Poutanen SM, Low DE, Jenkins SG, Katz K, Mulvey MR. 2011. Plasmid comparison and molecular analysis of *Klebsiella pneumoniae* harbouring *bla*_{KPC} from New York City and Toronto. *J Antimicrob Chemother* 66:1273–1277. <http://dx.doi.org/10.1093/jac/dkr092>.
- Kitchel B, Rasheed JK, Endimiani A, Hujer AM, Anderson KF, Bonomo RA, Patel JB. 2010. Genetic factors associated with elevated carbapenem resistance in KPC-producing *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* 54:4201–4207. <http://dx.doi.org/10.1128/AAC.00008-10>.
- Deleo FR, Chen L, Porcella SF, Martens CA, Kobayashi SD, Porter AR, Chavda KD, Jacobs MR, Mathema B, Olsen RJ, Bonomo RA, Musser JM, Kreiswirth BN. 2014. Molecular dissection of the evolution of carbapenem-resistant multilocus sequence type 258 *Klebsiella pneumoniae*. *Proc Natl Acad Sci U S A* 111:4988–4993. <http://dx.doi.org/10.1073/pnas.1321364111>.
- Correia M, Boavida F, Grosso F, Salgado MJ, Lito LM, Cristino JM, Mendo S, Duarte A. 2003. Molecular characterization of a new class 3 integron in *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* 47:2838–2843. <http://dx.doi.org/10.1128/AAC.47.9.2838-2843.2003>.
- Poirel L, Carattoli A, Bernabeu S, Bruderer T, Frei R, Nordmann P. 2010. A novel IncQ plasmid type harbouring a class 3 integron from *Escherichia coli*. *J Antimicrob Chemother* 65:1594–1598. <http://dx.doi.org/10.1093/jac/dkq166>.
- Shibata N, Doi Y, Yamane K, Yagi T, Kurokawa H, Shibayama K, Kato H, Kai K, Arakawa Y. 2003. PCR typing of genetic determinants for metallo-β-lactamases and integrases carried by Gram-negative bacteria isolated in Japan, with focus on the class 3 integron. *J Clin Microbiol* 41:5407–5413. <http://dx.doi.org/10.1128/JCM.41.12.5407-5413.2003>.
- Laraki N, Galleni M, Thamm I, Riccio ML, Amicosante G, Frère J-M, Rossolini GM. 1999. Structure of In31, a *bla*_{TIMP}-containing *Pseudomonas aeruginosa* integron phylogenically related to In5, which carries an unusual array of gene cassettes. *Antimicrob Agents Chemother* 43:890–901.

20. Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. 2005. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 63:219–228. <http://dx.doi.org/10.1016/j.mimet.2005.03.018>.
21. Cantón R, Akóva M, Carmeli Y, Giske CG, Glupczynski Y, Gniadkowski M, Livermore DM, Miriagou V, Naas T, Rossolini GM, Samuelsen Ø, Seifert H, Woodford N, Nordmann P, European Network on Carbapenemases. 2012. Rapid evolution and spread of carbapenemases among *Enterobacteriaceae* in Europe. *Clin Microbiol Infect* 18:413–431. <http://dx.doi.org/10.1111/j.1469-0691.2012.03821.x>.
22. Giske CG, Fröding I, Hasan CM, Turlej-Rogacka A, Toleman M, Livermore D, Woodford N, Walsh TR. 2012. Diverse sequence types of *Klebsiella pneumoniae* contribute to the dissemination of *bla*_{NDM-1} in India, Sweden, and the United Kingdom. *Antimicrob Agents Chemother* 56:2735–2738. <http://dx.doi.org/10.1128/AAC.06142-11>.
23. Woodford N, Turton JF, Livermore DM. 2011. Multiresistant Gram-negative bacteria: the role of high-risk clones in the dissemination of antibiotic resistance. *FEMS Microbiol Rev* 35:736–755. <http://dx.doi.org/10.1111/j.1574-6976.2011.00268.x>.