

Review Article

Posttranscriptional Regulation of Splicing Factor SRSF1 and Its Role in Cancer Cell Biology

Vânia Gonçalves^{1,2} and Peter Jordan^{1,2}

¹Department of Human Genetics, National Health Institute Dr. Ricardo Jorge, Avenida Padre Cruz, 1649-016 Lisboa, Portugal

²Biosystems and Integrative Sciences Institute (BioISI), Faculty of Sciences, University of Lisbon, 1749-016 Lisboa, Portugal

Correspondence should be addressed to Peter Jordan; peter.jordan@insa.min-saude.pt

Received 20 October 2014; Accepted 16 December 2014

Academic Editor: Elena Orlova

Copyright © 2015 V. Gonçalves and P. Jordan. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Over the past decade, alternative splicing has been progressively recognized as a major mechanism regulating gene expression patterns in different tissues and disease states through the generation of multiple mRNAs from the same gene transcript. This process requires the joining of selected exons or usage of different pairs of splice sites and is regulated by gene-specific combinations of RNA-binding proteins. One archetypical splicing regulator is SRSF1, for which we review the molecular mechanisms and posttranscriptional modifications involved in its life cycle. These include alternative splicing of SRSF1 itself, regulatory protein phosphorylation events, and the role of nuclear versus cytoplasmic SRSF1 localization. In addition, we resume current knowledge on deregulated SRSF1 expression in tumors and describe SRSF1-regulated alternative transcripts with functional consequences for cancer cell biology at different stages of tumor development.

1. Introduction

The expression of a gene is initiated by its transcription into a precursor messenger RNA (mRNA), which is then further processed and spliced into a mature mRNA. Splicing is regulated through the interaction between RNA-binding proteins (RBPs) and their cognate splicing regulatory sequence elements (SREs) in the mRNA. This is especially important for alternative splicing where multiple mRNAs can be generated from the same pre-mRNA through the joining of selected exons or usage of different pairs of splice sites [1].

The number of genes encoding RBPs in the human genome is currently estimated to be around 860 [2, 3], far below the number of around 200 000 transcripts that can be produced from the roughly 21 000 human protein-coding genes. Therefore, a key principle in splicing regulation is that the interaction of RNA-binding proteins with SREs is not a one-to-one relationship: each SRE motif can be recognized by multiple alternative RBPs and most splicing factors can recognize two or more SRE motifs. This is particularly relevant for alternative splicing events, the regulation of which involves a complex network of competing protein-RNA interactions

so that individual exons can be controlled by multiple factors [4, 5]. For example, multiple RNA-binding proteins with similar splicing regulatory activities might bind the same motif and this functional redundancy creates robustness in a splicing decision. Also, some factors compete with or displace another factor with opposite activity and confer functional antagonism. These overlapping binding specificities allow regulatory plasticity, which underlies tissue-specific splicing patterns, subtle fine-tuning of splice variant levels, and regulatory relationships between splicing regulators and upstream signaling pathways.

Among the RBPs, the major classes of splicing factors that control splice site recognition are the families of Serine/Arginine-rich (SR) proteins and heterogeneous nuclear ribonucleoproteins (hnRNPs). These proteins act by selecting splice sites for recognition by the spliceosome through binding to intronic or exonic splice enhancer and silencer elements and promoting or destabilizing protein interactions with spliceosome components. One of the best studied factors is SRSF1, formerly known as ASF or SF2 [6]. SRSF1 is a prototypical splicing factor mostly recruited to SREs classified as exonic splicing enhancers (ESEs). SRSF1

recognizes degenerate purine-rich sequence motifs [7, 8] and its binding promotes recognition of both constitutive and alternative exons during spliceosomal assembly. The current knowledge about its regulation will be the focus of this review. The described principles of regulation also apply to many other SR proteins and RBPs.

2. Posttranscriptional Regulation of SRSF1 mRNA

The SRSF1 gene is essential for normal embryonic development that is constitutively expressed and tightly regulated at the posttranscriptional level. In particular, SRSF1 recognizes SREs in its own transcripts, leading to alternative splicing, with some transcript forms being degraded by nonsense-mediated mRNA decay (NMD). In case of SRSF1, alternative splicing occurs in the 3' untranslated region following excision of an additional intron and thus introduction of a new exon-exon junction. In consequence, the original stop codon is recognized as premature and the transcript targeted for NMD [9]. This mechanism is highly conserved and shared by other SR proteins. It serves both as a negative feedback loop, in which increased SR protein levels promote an increase in unproductive splice variants of their own transcripts, and as a target for regulation, for example, depending on the ERK1/2-mediated phosphorylation status of the splicing regulator Sam68 [10].

In addition, other posttranscriptional mechanisms of SRSF1 autoregulation were described such as nuclear retention of alternative SRSF1 transcript variants or regulation of the translational efficiency of its transcripts [11, 12]. Furthermore, miRNAs targeting SRSF1 translation have begun to be identified, including miR-28, miR-505, miR-10a, and miR-10b [13, 14]. Thus, SRSF1 transcript levels are fine-tuned by various posttranscriptional mechanisms but the quantitative contribution of each step and their orchestration in response to different cellular stimuli remains undetermined.

3. SRSF1 Regulation by Protein Phosphorylation

Following translation of SRSF1 transcripts into protein, constitutive phosphorylation steps occur. First, the predominantly cytoplasmic SR-specific protein kinases (SRPKs) phosphorylate part of the C-terminal Arg-Ser-rich (RS) domain, which contains 20 serine residues. SRPK1 was shown to phosphorylate the proximal first 12 residues [15] and this promotes nuclear import through interaction of phospho-SRSF1 with the import factor transportin-SR2 [16] and subsequent localization into nuclear speckles [17, 18].

Once in the nucleus the Cdc2-like kinases (CLKs) phosphorylate the remaining serine residues in the distal RS domain which leads to dispersed nuclear localization of SRSF1 and is required for its function in splicing [19–21] through cotranscriptional association with RNA polymerase II (pol2). Upon transcription inhibition SRSF1 is translocated from the nucleoplasm back to nuclear speckles [22] (see Figure 1 for a graphic summary).

Recently, it was found that SRPK1 can also shuttle into the nucleus [23–26] where then SRPK1 and CLK1 display similar activities toward Arg-Ser repeats in the distal RS domain, suggesting that these kinases no longer operate in a strict linear manner along the RS domain. Instead, CLKs appear to recognize preferentially the three Ser-Pro dipeptides in the RS domain, the phosphorylation of which has been proposed to change the conformation of the RS domain and regulate SRSF1 contact sites required in the spliceosome [27]. Nuclear translocation of SRPK1 can be induced, for example, by stress conditions, and involves disruption of its binding to a cytoplasmic Hsp70/Hsp90 complex [28].

Besides these phosphorylation events considered to be constitutive, other protein kinases have been reported to regulate SRSF1 through phosphorylation.

Protein kinase A (PKA) can phosphorylate SRSF1 on serine 119 *in vitro* and modulate its activity as a splicing factor [29, 30]. This phosphorylation occurs in the so-called pseudo-RNA recognition motif (RRM) and was described to change the RNA-binding properties of SRSF1 and reduce its capacity to activate splicing.

Human DNA topoisomerase I (topo I) has also been described to phosphorylate SRSF1 [31, 32], most likely in the RS domain. This phosphorylation promotes the binding of SRSF1 to cognate ESEs during alternative splicing events [33]. The DNA damage signal poly-ADP ribose forms a complex with SRSF1, and this promotes Topo I to switch from its protein kinase to DNA relaxation activity [34]. A further connection between SRSF1 and Topo I is their role in preventing R-loop formation, stable mRNA:DNA hybrids that can form following transcription [35, 36].

Another kinase reported to phosphorylate SRSF1 *in vitro* is AKT [37, 38], which also targets serine residues in the RS domain, leading to altered splicing decisions. A subsequent study reported that AKT1 interacts with and promotes SRPK1 and SRPK2 autophosphorylation and their subsequent translocation into the nucleus [26] with simultaneous formation of a phosphatase containing complex to downregulate AKT activity [39]. It remains to be established whether AKT phosphorylates SRSF1 directly or whether the described activity of immunoprecipitated AKT to phosphorylate SRSF1 originates from associated SRPKs.

The serine/threonine kinase NEK2 is also a splicing factor kinase that colocalizes with SRSF1 in nuclear speckles. It interacts with and phosphorylates SRSF1, affecting the splicing activity of SRSF1 in a SRPK1-independent manner [40].

4. SRSF1 Regulation through Nuclear-Cytoplasmic Distribution

Ample experimental evidence showed that SRSF1 is a shuttling protein that localizes to both the nucleus and the cytoplasm, depending on the phosphorylation state of its RS domain [37, 41]. Furthermore, experimental blocking of SRPK (by either depleting its expression level or inhibiting its kinase activity) revealed that the cytoplasmic pool of SRSF1 increased, identifying phosphorylation as a major factor for




FIGURE 1: Life cycle and posttranslational modifications of the SRSF1 protein. Following its *de novo* synthesis at ribosomes, the cytoplasmic SRSF1 protein is constitutively phosphorylated by the cytosolic protein kinase SRPK at serine residues in the proximal RS domain (pRS). This first phosphorylation step is required for nuclear import of SRSF1, followed by a second phosphorylation step at the distal RS domain (dRS), including several serine-proline (SP) motifs. This step is usually catalyzed by the nuclear protein kinase CLK1 but can also be performed by SRPK if induced to translocate into the nucleus. Nuclear translocation is further modulated through methylation of the three arginine residues R93, R97, and R109 located between the two RNA recognition motif (RRM) domains (not shown). SRSF1 with a completely phosphorylated RS domain accumulates in nuclear speckles from where it is recruited to the spliceosome. SRSF1 dephosphorylation induces its nuclear to cytoplasmic translocation and lack of phosphorylation by SRPK in the cytosol leads to its proteolytic degradation.

SRSF1 nuclear translocation [42, 43]. The contribution of nuclear phosphatase activity to cytoplasmic export of SRSF1 has not been directly demonstrated *in vivo* but protein phosphatase 1 can dephosphorylate the proximal RS domain of SRSF1 *in vitro* or in permeabilized cell nuclei [44–46].

One physiological condition modulating SRSF1 localization is stress response, when general splicing is inhibited but specific alternative splicing events continue to occur [47]. For example, replicative senescence or induced stress stimuli of the vascular endothelium result in preferential cytoplasmic localization of SRSF1 and the underlying mechanism was postulated to involve nuclear import of SRPK1 and consequently lack of constitutive cytosolic SRSF1 phosphorylation [48]. In contrast, hyperphosphorylation of SRSF1 was observed during the DNA damage response and caused altered subnuclear distribution and changes in alternative splicing pattern of target genes [49].

Another posttranslational modification involved in SRSF1 localization is the methylation of three arginine residues (R93, R97, and R109) located in a region between the two RRM domains [50]. Lack of methylation in a triple-Ala mutant turned SRSF1 predominantly cytoplasmic, whereas

a triple-Lys substitution maintaining the positive charge localized to nuclear speckles, as the wild-type protein. How the respective protein arginine methyltransferases (PRMTs) are regulated and contribute to the nuclear-cytoplasmic transitions of SRSF1 is poorly understood.

Once in the nucleus, the long noncoding RNA MALAT1 (metastasis-associated lung adenocarcinoma transcript 1) interacts with SRSF1, which is important for the recruitment of other SR proteins into nuclear speckles [51].

5. Functional Consequences of Nuclear versus Cytoplasmic SRSF1 Concentrations

Relative concentrations of antagonizing or competing SFs are important determinants in alternative splicing regulation. For example, SRSF1 generally displays a stimulatory role in splicing when bound to exons and its function in alternative splicing *in vitro* can be antagonized by the activity of hnRNP A proteins in a concentration-dependent manner. *In vivo*, competition between SFs can originate from the relative ratios of such antagonists expressed in different tissues or developmental stages, creating tissue or stage-specific patterns of

splicing. In addition, the dynamic regulation of subcellular SF localization allows cells to modulate the effective nuclear concentration of a given SF and alter the pattern of expressed splicing variants in response to external stimuli. For example, the subnuclear distribution of SRSF1 changes during the DNA damage response following hyperphosphorylation and results in a shift in the alternative splicing pattern of target genes that control cell survival [49]. Also, drug-induced disruption of nuclear speckles with concomitant release of SRSF1 into the nucleoplasm induced changes in alternative splicing events [52]. And endothelial senescence is associated with a scattered distribution of SRSF1 throughout the cytoplasm. This leads to the expression of alternative isoforms of target genes such as endoglin (ENG), vascular endothelial growth factor A (VEGFA), tissue factor (T3), or lamin A (LMNA) that integrate into a common molecular senescence program [48]. Vice versa, epithelial cells treated with insulin-like growth factor-1 (IGF-1) displayed nuclear translocation of SRSF1, which was dependent on SPRK1/2 activity.

6. Regulation of SRSF1 by Cytosolic Protein Degradation

A specific decrease in SRSF1 protein levels was observed in SRPK1-depleted or SRPIN340-treated colorectal cells, without changes in the corresponding SRSF1 mRNA. This suggests that cytoplasmic SRSF1 localization leads to protein degradation. Indeed, the SRSF1 protein remained stable in such treated cells when incubated with inhibitors MG132 or lactacystin, indicating degradation by the proteasome [43]. It should be noted that studying SFs with proteasome inhibitors needs to be well controlled at the corresponding transcript level because the inhibitors are likely to affect other SFs or transcription factors in the cell. For example, the gene encoding SRSF3 (former SRp20) is a direct transcriptional target of β -catenin/TCF4 [53] so that inhibition of β -catenin degradation will increase expression of SRSF3, which in turn can promote unproductive alternative SRSF1 transcripts [54].

SRSF1 protein expression levels did also not correlate with mRNA expression levels following T cell stimulation. Immunoprecipitation studies showed increased ubiquitylation of SRSF1 in activated T cells and proteasomal but not lysosomal degradation was shown to be involved by blocking with specific inhibitors MG132 and bafilomycin, respectively. Interestingly, T cells from patients with SLE (systemic lupus erythematosus) showed increased ubiquitylation of SRSF1 when compared to those from healthy individuals [55].

Downregulation of SRSF1 protein level was further found to occur following inhibition of activity or siRNA-mediated depletion of GSK3 β in U87 or U373 glioblastoma cells [56]. Similarly, GSK3 β depletion in HT29 colorectal cancer cells led to a reduction in both SRSF1 and SRPK1 protein levels, suggesting an indirect effect of GSK3 β on SRSF1 via SRPK1 [43].

It has been described that the RS domain, which is common to all SR proteins, is required for their proteolytic degradation by the proteasome [57] but further mechanistic details remain to be determined.

7. Impact of Posttranslational Modification on Other RNA-Related Functions of SRSF1

SRSF1 has been shown to facilitate the nuclear export of spliced mRNAs to which it is bound through its interaction with the TAP/NXF1 receptor [58]. Interestingly, this adaptor function implies partial dephosphorylation of its RS domain for cytoplasmic translocation [59], suggesting the phosphorylation status of SRSF1 serves to regulate nuclear export of some mRNPs.

The subsequent ribosomal translation of transcripts containing a SRSF1-targeted ESE is also stimulated, both *in vivo* and *in vitro* [60]. Thus, SRSF1-mediated alternative splicing, mRNA export, and translational efficiency of its target transcripts are coupled, and their number has been identified experimentally to be around 500 [61].

In addition, SRSF1 overexpression was found to increase the ratio between cap-dependent and internal ribosome entry site-dependent translation initiation [62], probably by suppressing the activity of 4E-BP, a competitive inhibitor of cap-dependent translation [63]. Probably related to these properties is the observation that SRSF1 enhances nonsense-mediated mRNA decay (NMD) [64] because SRSF1 overexpression can promote the pioneer round of translation required for NMD to occur [58].

Another class of RNA pol2 transcripts is miRNAs and SRSF1 overexpression in HeLa cells promoted the maturation step of miR-7 and other miRNAs. SRSF1 (and also other splicing factors) directly interacts with primary miR transcripts and promotes the Drosha cleavage step generating mature miRNA [65].

Another recently discovered function of SRSF1 is to enhance protein sumoylation [66]. SRSF1 associates with the SUMO E2 conjugating enzyme Ubc9 and enhances SUMO conjugation to RNA processing factors but further details on the regulation or consequences of this modification remain to be identified.

It should also be noted that SRSF1 was shown to be involved in chromatin organization and histone modifications such as H3K36me3, which are relevant for splicing decisions [67, 68].

8. Role of SRSF1 in Cancer Cell Biology and Tumorigenesis

Malignant changes in the cellular genome can either be tumor-initiating driver events or subsequent adaptations required for tumor cell progression. Such changes either alter the expression level of critical genes or their nucleotide sequence to generate gain- or loss-of-function mutant gene products. Although point mutations in core components of the spliceosome were recently discovered using whole-genome sequencing approaches [69], reports from various tumor types revealed that splicing factors mostly show increased expression levels [70–72]. Concerning SRSF1, overexpression was reported in tumors from colon, thyroid, small intestine, kidney, lung, liver, pancreas, and breast [73, 74]. In childhood acute lymphoblastic leukemia SRSF1

was further found to be upregulated together with protein arginine methyltransferase PRMT1 [75], which is involved in promoting SRSF1 nuclear localization [50].

The overexpression of SRSF1 in tumors has been related to several alternative mechanisms. First, in breast tumors and breast cancer cell lines amplification of the *SFRS1* gene at chromosomal location 17q23 was detected and the increased DNA copy number correlated with elevated SRSF1 mRNA levels [73].

Second, the *SRSF1* gene is a target of MYC, a potent oncogenic transcription factor overexpressed in many different tumor types that has pleiotropic effects on cancer cell biology [76]. MYC binds directly to the SRSF1 promoter and activates transcription. Both genes were found coexpressed in lung and breast carcinomas and MYC depletion downregulates SRSF1 expression in lung cancer cell lines [77].

Third, the above-mentioned negative feedback loop, in which SRSF1 promotes an increase in unproductive splice variants of its own transcripts, can be subverted in the presence of splicing regulator SAM68. Changes in expression or phosphorylation of SAM68 were found to promote the formation of full-length SRSF1 transcripts, thus leading to increased SRSF1 protein levels [10]. SAM68 phosphorylation depends on ERK/MAP kinase activity, which is frequently augmented in human tumors.

Together, this indicates that SRSF1 overexpression is in general the consequence of other preceding tumor-initiating genetic changes but contributes to further tumor progression.

Two apparently opposing consequences of SRSF1 overexpression on cancer cell biology have been described: the induction of oncogene-induced senescence and the malignant transformation of cells. On the one hand, SRSF1 overexpression leads to the formation of a nucleoplasmic complex with the ribosomal protein RPL5 and the E3-ubiquitin ligase MDM2, which normally ubiquitylates the p53 tumor-suppressor protein leading to its proteolytic degradation [78]. Complex formation inhibits MDM2 and thus p53 protein levels increase and trigger a cellular senescence response, which normally is part of a ribosomal stress pathway. Because the ability of SRSF1 overexpression to activate a tumor-suppressing senescence response is dependent on an intact p53 pathway, the identified SRSF1-overexpressing tumor types revealed characteristics of p53 inactivation [78].

On the other hand, SRSF1 can act as an oncogene since a twofold increase in expression can transform immortalized rodent fibroblasts [73, 79] and human mammary epithelial cells [73, 79]. In these models, SRSF1 overexpression promoted cell proliferation and antiapoptotic pathways, mainly reflecting the combined effects of several alternative splicing variants which were activated by the concentration-dependent changes in SRSF1 availability. Some of these specific variants have been characterized, as detailed below, but probably represent just the tip of the iceberg.

One group of identified target genes is formed by the apoptosis regulators BIN1, BCL2L1 (BIM), BCL-XL, ICAD, and MCL1, with SRSF1 overexpression in cancer cells promoting the formation of their respective antiapoptotic splice variants. Several target genes belong to the Bcl-2 family of proteins, which regulate whether the Bak and Bax proteins

can cause mitochondrial outer membrane permeabilization and cytochrome c release as the trigger for intrinsic apoptosis induction. The Bcl-2 family comprises both proapoptotic and antiapoptotic proteins, depending on their BH domain composition, and it is the balance between both types of proteins that determines whether the mitochondrial pathway to apoptosis is activated [80].

Regarding BIM, several SRSF1-induced transcript variants were described lacking exons 2, 3, or 4 (BIM γ 1, γ 2, ES) which encode the BH3 domain. This domain binds antiapoptotic Bcl-2 family members and is necessary for induction of apoptosis by BIM [79, 81]. Similarly, SRSF1 expression promotes inclusion of exon 2 of the BH3 domain-containing gene MCL-1 (myeloid cell leukemia-1) giving rise to the antiapoptotic MCL-1L isoform in both breast cancer and choriocarcinoma cells [82]. Overexpression of SRSF1 also promotes generation of the antiapoptotic isoform BCL-XL [83].

BIN1 has tumor-suppressor activity by interacting with and activating MYC-mediated apoptosis, except when exon 12A is included by SRSF1-mediated alternative splicing, because the resulting antiapoptotic BIN1+12A isoform is unable to interact with MYC. Furthermore, SRSF1 was shown to modulate exclusion of exon 5 of the mRNA encoding the inhibitor of caspase-activated DNase (ICAD), a regulator of the DNase responsible for DNA fragmentation during apoptosis [84].

A parallel group of SRSF1-regulated target genes is involved in cellular signaling pathways related to proliferation and cell cycle progression. Examples of genes from this group are *CCND1*, *RPS6KBI*, *RON*, *RAC1*, and *MKNK2* genes.

SRSF1 increases expression of the cyclin D1b oncogene which arises from alternative splicing of the *CCND1* transcript, and harbors enhanced oncogenic functions not shared by full-length cyclin D1 (cyclin D1a) [85]. In this case, SRSF1 blocks recognition of the *CCND1* exon 4-intron 4 boundary, thus repressing inclusion of exon 5 so that a nuclear protein with a unique C-terminus is generated. SRSF1 also promotes the inclusion of exon 5 into the pre-mRNAs of *TEAD-1* (TEF-1 or TCF13), a transcription factor normally involved in cell differentiation and cell cycle arrest in myoblasts [86].

The *RPS6KBI* gene encodes the protein S6 kinase 1, a substrate for the cell growth regulating kinase mTOR. Excess SRSF1 promoted an increase in S6K1 variants by including one to three alternative cassette exons between exons 6 and 7 that are normally skipped [73] and include a proper stop codon. These short S6K1 isoforms have a truncated kinase domain and lack the mTOR-regulated C-terminus but are able to bind to and activate the mTORC1 complex. This activation of the mTORC1 complex occurs independent of the classical PI3K/AKT pathway and leads to phosphorylation of eIF4EBP1, releasing its inhibitory effect on cap-dependent translation [87, 88]. In addition, there is evidence that SRSF1 itself participates in a complex with mTORC1 to enhance translation efficiency of its target transcripts [63], for example, survivin [89] and β -catenin [90].

RON encodes a receptor tyrosine kinase in breast and colon tumors and SRSF1 promotes skipping of exon 11 by binding to an enhancer element in the competing exon 12.

The resulting isoform Δ Ron is constitutively active and promotes cell motility [91] as part of an epithelial-mesenchymal transition program, which tumor cells may use to escape from adverse local growth conditions.

Breast and colon tumors are further characterized by overexpression of Rac1b [92, 93], a hyperactivated splice variant of the small GTPase Rac1, which is involved in gene transcription and cell motility [94, 95]. In colorectal cells, SRSF1 was shown to be required for inclusion of an additional exon 3b to generate Rac1b [96] and increased expression of Rac1b contributes to cell survival [97, 98].

Finally, SRSF1 overexpression enhances the inclusion of the alternative 3'-terminal exon 13b of the gene *MKNK2* encoding the protein kinase Mnk2, an effector in the ERK/MAPK pathway [73]. The corresponding isoform Mnk2b lacks a C-terminal MAPK-binding domain and does not phosphorylate and activate the p38-MAPK required for stress-induced cell death. In contrast, it sustains phosphorylation of the translation initiation factor eIF4E, thus promoting cap-dependent protein translation and cell growth [99].

Curiously, splicing of the Mnk2b isoform was not induced when a chimeric nucleus-retained SRSF1 protein (SRSF1-NRS1), fused to the nuclear retention signal of the non-shuttling protein SRSF2, was overexpressed, suggesting an indirect effect of SRSF1 for this splicing event. Nevertheless, SRSF1-NRS1 was as competent as wild-type SRSF1 in inducing mammary cell transformation in 3D cultures but requires presence of the RRM1 domain, revealing a significant contribution of the nuclear functions of SRSF1 to cell transformation [79].

In hepatocellular xenografts, however, SRSF1-NRS1 protein had a much lower effect on tumor formation than SRSF1. In this model, SRSF1 overexpression also promotes activation of ERK/MAPK, probably by increasing B-RAF mRNA and protein levels. Although the mechanism remains to be explained, the RRM1 domain is required and sufficient to induce activation of this oncogenic pathway, indicating that the effect occurs at the mRNA level [81]. These findings reveal that SRSF1 can exert its oncogenic role through both nuclear and cytosolic pathways depending on the cellular contexts.

More recently, SRSF1 overexpression was also reported in lung cancer and novel SRSF1 target transcripts were identified, including the genes, *ATP11C*, *IQCB1*, *TUBD1*, proline-rich coiled-coil 2C (*PRRC2C*) [100], and survivin [89].

Apart from the genetic changes affecting proliferation and survival of the transformed cancer cells themselves, another important aspect of cancer cell biology is their cellular communication with the surrounding stroma. It is imperative for the growth of epithelial tumors to gain access to nutrient supply via blood vessels so that cancer cells release angiogenic signals to endothelial cells, for instance, the vascular endothelial growth factor VEGF. SRSF1 is involved in promoting proximal splice site selection in C-terminal exon 8 of VEGF, resulting in the generation of proangiogenic isoforms [101, 102]. Besides overexpression, this can result from SRSF1 activation in epithelial cells following oncogenic signaling through IGF-1, EGF, or TNF- α . These factors lead to activation of SRPKs [26, 42], which then phosphorylate SR proteins including SRSF1. SRPK1 inhibition has been used to

manipulate the local balance of pro- and antiangiogenic in eye pathologies caused by neovascularization [103] and might be interesting for cancer therapy. In addition, SRSF1-mediated alternative splicing of both Ron and TEAD-1 has been linked to increased expression of angiogenic growth factors [77, 104]. SRSF1 may thus impact VEGF expression through both direct and indirect regulation to promote angiogenesis.

The role of fibroblasts in the stroma is to deposit or remodel extracellular matrix components and this is important for tumor cell migration. For example, a dense fibronectin meshwork favors epithelial cell invasion and results from inclusion of the EDA exon through SRSF1-regulated alternative splicing of the unique fibronectin-encoding gene [105, 106]. This occurs during embryogenesis but also in adult fibroblasts during tissue repair, tumor progression, and inflammation when expression levels of SRSF1 increase. It remains to be established whether tumor cells can release signals that induce increased SRSF1 expression in tumor-associated fibroblasts.

Following therapeutic challenge of tumor cells with DNA-damaging agents, resistant cells can eventually emerge. In one report hyperphosphorylation of SRSF1 was observed in the presence of DNA damage, causing altered subnuclear distribution and changes in alternative splicing pattern of target genes that promote cell survival [49]. Similarly, treatment of pancreatic tumor cells with the nucleoside analogue gemcitabine induced SRSF1 overexpression, and the resulting splicing of *MNK2b* with consequent phosphorylation of the translation initiation factor eIF4E was identified as the cause for drug resistance [107]. Furthermore, non-small cell lung cancer (NSCLC) cells respond to daunorubicin or cisplatin with an antiapoptotic caspase 9b splice variant. SRSF1 regulates this alternative splicing event by binding to a splicing enhancer in intron 6 and subsequent exclusion of an exon 3,4,5,6-cassette, generating caspase 9b [108]. SRSF1 is activated following hyperphosphorylation at serines 199, 201, 227, and 234 [109], mediating the therapeutic resistance of NSCLC. Another study in NSCLC observed that SRSF1 protein accumulates when cells were treated with carboplatin and paclitaxel and that cells stably overexpressing SRSF1 were more resistant to these chemotherapeutic drugs [74].

9. Conclusions

SRSF1 is an important protein for the regulation of constitutive and alternative splicing of cellular pre-mRNAs. Its activity as splicing regulator depends on the relative expression level of SRSF1 compared to other antagonistic or synergistic splicing factors as well as on its posttranslational modifications. In particular, the phosphorylation state of SRSF1 determines its nuclear or cytoplasmic localization and proteolytic degradation. Overexpression of SRSF1 has been reported in various tumors types and this has consequences for the alternative splicing profile expressed in tumor cells. Clear experimental evidence for tumor-promoting effects of SRSF1-induced alternative splicing variants has been provided but the genome-wide scale of its effects on cancer cell biology remains to be described.

Similar studies on other splicing factors are beginning to emerge and will likely reveal comparable complex effects on cancer cell transcriptomes as part of an adaptive response to activate survival pathways in tumor cells. A more comprehensive knowledge of these pathways may allow designing therapeutic interventions based on a combination of inhibitory drugs targeting simultaneously various pathways to reduce the selection of therapy-resistant tumor cell clones.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

Recent work in the authors' lab was supported by the Fundação para a Ciência e Tecnologia, Portugal (Grants PEst-OE/BIA/UI4046/2011 to the BioFIG research unit, FCT/MCTES/PIDDAC to BioISI Research Centre UID/MULTI/04046/2013, and fellowship BPD 63395/2009 to Vânia Gonçalves). The authors thank Drs. Paulo Matos and Margarida Gama-Carvalho for their longstanding collaboration in studying alternative splicing.




References

- [1] M. Chen and J. L. Manley, "Mechanisms of alternative splicing regulation: insights from molecular and genomics approaches," *Nature Reviews Molecular Cell Biology*, vol. 10, no. 11, pp. 741–754, 2009.
- [2] A. Castello, B. Fischer, K. Eichelbaum et al., "Insights into RNA biology from an atlas of mammalian mRNA-binding proteins," *Cell*, vol. 149, no. 6, pp. 1393–1406, 2012.
- [3] D. Ray, H. Kazan, K. B. Cook et al., "A compendium of RNA-binding motifs for decoding gene regulation," *Nature*, vol. 499, no. 7457, pp. 172–177, 2013.
- [4] K. J. Hertel, "Combinatorial control of exon recognition," *The Journal of Biological Chemistry*, vol. 283, no. 3, pp. 1211–1215, 2008.
- [5] C. W. J. Smith and J. Valcárcel, "Alternative pre-mRNA splicing: the logic of combinatorial control," *Trends in Biochemical Sciences*, vol. 25, no. 8, pp. 381–388, 2000.
- [6] J. L. Manley and A. R. Krainer, "A rational nomenclature for serine/arginine-rich protein splicing factors (SR proteins)," *Genes and Development*, vol. 24, no. 11, pp. 1073–1074, 2010.
- [7] P. J. Smith, C. Zhang, J. Wang, S. L. Chew, M. Q. Zhang, and A. R. Krainer, "An increased specificity score matrix for the prediction of SF2/ASF-specific exonic splicing enhancers," *Human Molecular Genetics*, vol. 15, no. 16, pp. 2490–2508, 2006.
- [8] J. R. Sanford, X. Wang, M. Mort et al., "Splicing factor SFRS1 recognizes a functionally diverse landscape of RNA transcripts," *Genome Research*, vol. 19, no. 3, pp. 381–394, 2009.
- [9] L. F. Lareau, M. Inada, R. E. Green, J. C. Wengrod, and S. E. Brenner, "Unproductive splicing of SR genes associated with highly conserved and ultraconserved DNA elements," *Nature*, vol. 446, no. 7138, pp. 926–929, 2007.
- [10] C. Valacca, S. Bonomi, E. Buratti et al., "Sam68 regulates EMT through alternative splicing-activated nonsense-mediated mRNA decay of the SF2/ASF proto-oncogene," *The Journal of Cell Biology*, vol. 191, no. 1, pp. 87–99, 2010.
- [11] S. Sun, Z. Zhang, R. Sinha, R. Karni, and A. R. Krainer, "SF2/ASF autoregulation involves multiple layers of post-transcriptional and translational control," *Nature Structural and Molecular Biology*, vol. 17, no. 3, pp. 306–312, 2010.
- [12] J. R. Sanford, P. Coutinho, J. A. Hackett, X. Wang, W. Ranahan, and J. F. Cáceres, "Identification of nuclear and cytoplasmic mRNA targets for the shuttling protein SF2/ASF," *PLoS ONE*, vol. 3, no. 10, Article ID e3369, 2008.
- [13] L. Verduci, M. Simili, M. Rizzo et al., "MicroRNA (miRNA)-mediated interaction between leukemia/lymphoma-related factor (LRF) and alternative splicing factor/splicing factor 2 (ASF/SF2) affects mouse embryonic fibroblast senescence and apoptosis," *Journal of Biological Chemistry*, vol. 285, no. 50, pp. 39551–39563, 2010.
- [14] S. Meseguer, G. Mudduluru, J. M. Escamilla, H. Allgayer, and D. Baretino, "MicroRNAs-10a and -10b contribute to retinoic acid-induced differentiation of neuroblastoma cells and target the alternative splicing regulatory factor SFRS1 (SF2/ASF)," *The Journal of Biological Chemistry*, vol. 286, no. 6, pp. 4150–4164, 2011.
- [15] J. C. Hagopian, C.-T. Ma, B. R. Meade et al., "Adaptable molecular interactions guide phosphorylation of the SR protein ASF/SF2 by SRPK1," *Journal of Molecular Biology*, vol. 382, no. 4, pp. 894–909, 2008.
- [16] M.-C. Lai, R.-I. Lin, and W.-Y. Tarn, "Transportin-SR2 mediates nuclear import of phosphorylated SR proteins," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 98, no. 18, pp. 10154–10159, 2001.
- [17] J. F. Cáceres, T. Misteli, G. R. Sreaton, D. L. Spector, and A. R. Krainer, "Role of the modular domains of SR proteins in subnuclear localization and alternative splicing specificity," *Journal of Cell Biology*, vol. 138, no. 2, pp. 225–238, 1997.
- [18] N. Kataoka, J. L. Bachorik, and G. Dreyfuss, "Transportin-SR, a nuclear import receptor for SR proteins," *Journal of Cell Biology*, vol. 145, no. 6, pp. 1145–1152, 1999.
- [19] K. Colwill, T. Pawson, B. Andrews et al., "The Clk/Sty protein kinase phosphorylates SR splicing factors and regulates their intranuclear distribution," *EMBO Journal*, vol. 15, no. 2, pp. 265–275, 1996.
- [20] J. C. K. Ngo, S. Chakrabarti, J.-H. Ding et al., "Interplay between SRPK and Clk/Sty kinases in phosphorylation of the splicing factor ASF/SF2 is regulated by a docking motif in ASF/SF2," *Molecular Cell*, vol. 20, no. 1, pp. 77–89, 2005.
- [21] G. Ghosh and J. A. Adams, "Phosphorylation mechanism and structure of serine-arginine protein kinases," *The FEBS Journal*, vol. 278, no. 4, pp. 587–597, 2011.
- [22] T. Misteli, J. F. Cáceres, and D. L. Spector, "The dynamics of a pre-mRNA splicing factor in living cells," *Nature*, vol. 387, no. 6632, pp. 523–527, 1997.
- [23] J.-F. Gui, W. S. Lane, and X.-D. Fu, "A serine kinase regulates intracellular localization of splicing factors in the cell cycle," *Nature*, vol. 369, no. 6482, pp. 678–682, 1994.
- [24] J.-H. Ding, "Regulated cellular partitioning of SR protein-specific kinases in mammalian cells," *Molecular Biology of the Cell*, vol. 17, pp. 876–885, 2005.
- [25] T. Giannakouros, E. Nikolakaki, I. Mylonis, and E. Georgatsou, "Serine-arginine protein kinases: a small protein kinase family with a large cellular presence," *FEBS Journal*, vol. 278, no. 4, pp. 570–586, 2011.

- [26] Z. Zhou, J. Qiu, W. Liu et al., "The Akt-SRPK-SR axis constitutes a major pathway in transducing EGF signaling to regulate alternative splicing in the nucleus," *Molecular Cell*, vol. 47, no. 3, pp. 422–433, 2012.
- [27] B. E. Aubol, R. M. Plocinik, J. C. Hagopian et al., "Partitioning RS domain phosphorylation in an SR protein through the CLK and SRPK protein kinases," *Journal of Molecular Biology*, vol. 425, no. 16, pp. 2894–2909, 2013.
- [28] X.-Y. Zhong, J.-H. Ding, J. A. Adams, G. Ghosh, and X.-D. Fu, "Regulation of SR protein phosphorylation and alternative splicing by modulating kinetic interactions of SRPK1 with molecular chaperones," *Genes and Development*, vol. 23, no. 4, pp. 482–495, 2009.
- [29] J. Shi, W. Qian, X. Yin et al., "Cyclic AMP-dependent protein kinase regulates the alternative splicing of tau exon 10: a mechanism involved in tau pathology of Alzheimer disease," *The Journal of Biological Chemistry*, vol. 286, no. 16, pp. 14639–14648, 2011.
- [30] A. K. Aksaas, S. Eikvar, G. Akusjärvi, B. S. Skålhegg, and A. K. Kvissel, "Protein Kinase a-dependent phosphorylation of serine 119 in the proto-oncogenic serine/arginine-rich splicing factor 1 modulates its activity as a splicing enhancer protein," *Genes and Cancer*, vol. 2, no. 8, pp. 841–851, 2011.
- [31] F. Rossi, E. Labourier, T. Forné et al., "Specific phosphorylation of SR proteins by mammalian DNA topoisomerase I," *Nature*, vol. 381, no. 6577, pp. 80–82, 1996.
- [32] E. Labourier, F. Rossi, I.-E. Gallouzi, E. Allemand, G. Divita, and J. Tazi, "Interaction between the N-terminal domain of human DNA topoisomerase I and the arginine-serine domain of its substrate determines phosphorylation of SF2/ASF splicing factor," *Nucleic Acids Research*, vol. 26, no. 12, pp. 2955–2962, 1998.
- [33] J. Soret, M. Gabut, C. Dupon et al., "Altered serine/arginine-rich protein phosphorylation and exonic enhancer-dependent splicing in Mammalian cells lacking topoisomerase I," *Cancer Research*, vol. 63, no. 23, pp. 8203–8211, 2003.
- [34] M. Malanga, A. Czuby, A. Girstun, K. Staron, and F. R. Althaus, "Poly(ADP-ribose) binds to the splicing factor ASF/SF2 and regulates its phosphorylation by DNA topoisomerase I," *Journal of Biological Chemistry*, vol. 283, no. 29, pp. 19991–19998, 2008.
- [35] X. Li and J. L. Manley, "Inactivation of the SR protein splicing factor ASF/SF2 results in genomic instability," *Cell*, vol. 122, no. 3, pp. 365–378, 2005.
- [36] A. El Hage, S. L. French, A. L. Beyer, and D. Tollervey, "Loss of Topoisomerase I leads to R-loop-mediated transcriptional blocks during ribosomal RNA synthesis," *Genes and Development*, vol. 24, no. 14, pp. 1546–1558, 2010.
- [37] M. Blaustein, F. Pelisch, T. Tanos et al., "Concerted regulation of nuclear and cytoplasmic activities of SR proteins by AKT," *Nature Structural & Molecular Biology*, vol. 12, no. 12, pp. 1037–1044, 2005.
- [38] K. Jiang, N. A. Patel, J. E. Watson et al., "Akt2 regulation of Cdc2-like kinases (Clk/Sty), serine/arginine-rich (SR) protein phosphorylation, and insulin-induced alternative splicing of PKC β III messenger ribonucleic acid," *Endocrinology*, vol. 150, no. 5, pp. 2087–2097, 2009.
- [39] P. Wang, Z. Zhou, A. Hu et al., "Both decreased and increased SRPK1 levels promote cancer by interfering with PHLPP-mediated dephosphorylation of Akt," *Molecular Cell*, vol. 54, no. 3, pp. 378–391, 2014.
- [40] C. Naro, F. Barbagallo, P. Chieffi, C. F. Bourgeois, M. P. Paronetto, and C. Sette, "The centrosomal kinase NEK2 is a novel splicing factor kinase involved in cell survival," *Nucleic Acids Research*, vol. 42, no. 5, pp. 3218–3227, 2014.
- [41] D. Cazalla, J. Zhu, L. Manche, E. Huber, A. R. Krainer, and J. F. Cáceres, "Nuclear export and retention signals in the RS domain of SR proteins," *Molecular and Cellular Biology*, vol. 22, no. 19, pp. 6871–6882, 2002.
- [42] D. G. Nowak, E. M. Amin, E. S. Rennel et al., "Regulation of Vascular Endothelial Growth Factor (VEGF) splicing from pro-angiogenic to anti-angiogenic isoforms: a novel therapeutic strategy for angiogenesis," *Journal of Biological Chemistry*, vol. 285, no. 8, pp. 5532–5540, 2010.
- [43] V. Goncalves, A. Henriques, J. Pereira et al., "Phosphorylation of SRSF1 by SRPK1 regulates alternative splicing of tumor-related Rac1b in colorectal cells," *RNA*, vol. 20, no. 4, pp. 474–482, 2014.
- [44] T. Misteli and D. L. Spector, "Serine/threonine phosphatase 1 modulates the subnuclear distribution of pre-mRNA splicing factors," *Molecular Biology of the Cell*, vol. 7, no. 10, pp. 1559–1572, 1996.
- [45] T. Novoyatleva, B. Heinrich, Y. Tang et al., "Protein phosphatase 1 binds to the RNA recognition motif of several splicing factors and regulates alternative pre-mRNA processing," *Human Molecular Genetics*, vol. 17, no. 1, pp. 52–70, 2008.
- [46] C.-T. Ma, G. Ghosh, X.-D. Fu, and J. A. Adams, "Mechanism of dephosphorylation of the SR protein ASF/SF2 by protein phosphatase 1," *Journal of Molecular Biology*, vol. 403, no. 3, pp. 386–404, 2010.
- [47] G. Biamonti and J. F. Cáceres, "Cellular stress and RNA splicing," *Trends in Biochemical Sciences*, vol. 34, no. 3, pp. 146–153, 2009.
- [48] F. J. Blanco and C. Bernabéu, "The splicing factor SRSF1 as a marker for endothelial senescence," *Frontiers in Physiology*, vol. 3, article 54, 2012.
- [49] V. Leva, S. Giuliano, A. Bardoni et al., "Phosphorylation of SRSF1 is modulated by replicational stress," *Nucleic Acids Research*, vol. 40, no. 3, pp. 1106–1117, 2012.
- [50] R. Sinha, E. Allemand, Z. Zhang, R. Karni, M. P. Myers, and A. R. Krainer, "Arginine methylation controls the subcellular localization and functions of the oncoprotein splicing factor SF2/ASF," *Molecular and Cellular Biology*, vol. 30, no. 11, pp. 2762–2774, 2010.
- [51] V. Tripathi, D. Y. Song, X. Zong et al., "SRSF1 regulates the assembly of pre-mRNA processing factors in nuclear speckles," *Molecular Biology of the Cell*, vol. 23, no. 18, pp. 3694–3706, 2012.
- [52] Y. Kurogi, Y. Matsuo, Y. Mihara et al., "Identification of a chemical inhibitor for nuclear speckle formation: Implications for the function of nuclear speckles in regulation of alternative pre-mRNA splicing," *Biochemical and Biophysical Research Communications*, vol. 446, no. 1, pp. 119–124, 2014.
- [53] V. Gonçalves, P. Matos, and P. Jordan, "The β -catenin/TCF4 pathway modifies alternative splicing through modulation of SRp20 expression," *RNA*, vol. 14, no. 12, pp. 2538–2549, 2008.
- [54] Ú. Muñoz, J. E. Puche, R. Hannivoort, U. E. Lang, M. Cohen-Naftaly, and S. L. Friedman, "Hepatocyte growth factor enhances alternative splicing of the Krüppel-like factor 6 (KLF6) tumor suppressor to promote growth through SRSF1," *Molecular Cancer Research*, vol. 10, no. 9, pp. 1216–1227, 2012.
- [55] V. R. Moulton, A. R. Gillooly, and G. C. Tsokos, "Ubiquitination regulates expression of the serine/arginine-rich splicing factor 1 (SRSF1) in normal and systemic lupus erythematosus (SLE) T

- cells," *Journal of Biological Chemistry*, vol. 289, no. 7, pp. 4126–4134, 2014.
- [56] A. K. Yadav, V. Vashishta, N. Joshi, and P. Taneja, "AR-A 014418 used against GSK3beta downregulates expression of hnRNPA1 and SF2/ASF splicing factors," *Journal of Oncology*, vol. 2014, Article ID 695325, 9 pages, 2014.
- [57] O. Breig and F. Baklouti, "Proteasome-mediated proteolysis of SRSF5 splicing factor intriguingly co-occurs with SRSF5 mRNA upregulation during late erythroid differentiation," *PLoS ONE*, vol. 8, no. 3, Article ID e59137, 2013.
- [58] H. Sato, N. Hosoda, and L. E. Maquat, "Efficiency of the pioneer round of translation affects the cellular site of nonsense-mediated mRNA decay," *Molecular Cell*, vol. 29, no. 2, pp. 255–262, 2008.
- [59] M.-C. Lai and W.-Y. Tarn, "Hypophosphorylated ASF/SF2 binds TAP and is present in messenger ribonucleoproteins," *The Journal of Biological Chemistry*, vol. 279, no. 30, pp. 31745–31749, 2004.
- [60] J. R. Sanford, N. K. Gray, K. Beckmann, and J. F. Cáceres, "A novel role for shuttling SR proteins in mRNA translation," *Genes & Development*, vol. 18, no. 7, pp. 755–768, 2004.
- [61] M. M. Maslon, S. R. Heras, N. Bellora, E. Eyra, and J. F. Cáceres, "The translational landscape of the splicing factor SRSF1 and its role in mitosis," *eLife*, vol. 2014, no. 3, Article ID e02028, 2014.
- [62] M. Blaustein, L. Quadrona, G. Risso, M. De La Mata, F. Pelisch, and A. Srebrow, "SF2/ASF regulates proteomic diversity by affecting the balance between translation initiation mechanisms," *Journal of Cellular Biochemistry*, vol. 107, no. 4, pp. 826–833, 2009.
- [63] G. Michlewski, J. R. Sanford, and J. F. Cáceres, "The splicing factor SF2/ASF regulates translation initiation by enhancing phosphorylation of 4E-BP1," *Molecular Cell*, vol. 30, no. 2, pp. 179–189, 2008.
- [64] Z. Zhang and A. R. Krainer, "Involvement of SR proteins in mRNA surveillance," *Molecular Cell*, vol. 16, no. 4, pp. 597–607, 2004.
- [65] H. Wu, S. Sun, K. Tu et al., "A splicing-independent function of SF2/ASF in microRNA processing," *Molecular Cell*, vol. 38, no. 1, pp. 67–77, 2010.
- [66] F. Pelisch, J. Gerez, J. Druker et al., "The serine/arginine-rich protein SF2/ASF regulates protein sumoylation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 37, pp. 16119–16124, 2010.
- [67] R. J. Loomis, Y. Naoe, J. B. Parker et al., "Chromatin binding of SRp20 and ASF/SF2 and dissociation from mitotic chromosomes is modulated by histone H3 serine 10 phosphorylation," *Molecular Cell*, vol. 33, no. 4, pp. 450–461, 2009.
- [68] M. M. Pradeepa, H. G. Sutherland, J. Ule, G. R. Grimes, and W. A. Bickmore, "Psp1/Ledgf p52 binds methylated histone H3K36 and splicing factors and contributes to the regulation of alternative splicing," *PLoS Genetics*, vol. 8, no. 5, Article ID e1002717, 2012.
- [69] V. Visconte, H. Makishima, J. P. MacIejewski, and R. V. Tiu, "Emerging roles of the spliceosomal machinery in myelodysplastic syndromes and other hematological disorders," *Leukemia*, vol. 26, no. 12, pp. 2447–2454, 2012.
- [70] C. Ghigna, M. Moroni, C. Porta, S. Riva, and G. Biamonti, "Altered expression of heterogeneous nuclear ribonucleoproteins and SR factors in human colon adenocarcinomas," *Cancer Research*, vol. 58, no. 24, pp. 5818–5824, 1998.
- [71] E. Stickeler, F. Kittrell, D. Medina, and S. M. Berget, "Stage-specific changes in SR splicing factors and alternative splicing in mammary tumorigenesis," *Oncogene*, vol. 18, no. 24, pp. 3574–3582, 1999.
- [72] J. P. Venables, "Aberrant and alternative splicing in cancer," *Cancer Research*, vol. 64, no. 21, pp. 7647–7654, 2004.
- [73] R. Karni, E. De Stanchina, S. W. Lowe, R. Sinha, D. Mu, and A. R. Krainer, "The gene encoding the splicing factor SF2/ASF is a proto-oncogene," *Nature Structural and Molecular Biology*, vol. 14, no. 3, pp. 185–193, 2007.
- [74] S. Gout, E. Brambilla, A. Boudria et al., "Abnormal expression of the pre-mRNA splicing regulators SRSF1, SRSF2, SRPK1 and SRPK2 in non small cell lung carcinoma," *PLoS ONE*, vol. 7, no. 10, Article ID e46539, 2012.
- [75] L. Zou, H. Zhang, C. Du et al., "Correlation of SRSF1 and PRMT1 expression with clinical status of pediatric acute lymphoblastic leukemia," *Journal of Hematology and Oncology*, vol. 5, article 42, 2012.
- [76] W. P. Tansey, "Mammalian MYC proteins and cancer," *New Journal of Science*, vol. 2014, Article ID 757534, 27 pages, 2014.
- [77] S. Das, O. Anczuków, M. Akerman, and A. R. Krainer, "Oncogenic splicing factor SRSF1 is a critical transcriptional target of MYC," *Cell Reports*, vol. 1, no. 2, pp. 110–117, 2012.
- [78] O. I. Fregoso, S. Das, M. Akerman, and A. R. Krainer, "Splicing-factor oncoprotein SRSF1 stabilizes p53 via RPL5 and induces cellular senescence," *Molecular Cell*, vol. 50, no. 1, pp. 56–66, 2013.
- [79] O. Anczuków, A. Z. Rosenberg, M. Akerman et al., "The splicing factor SRSF1 regulates apoptosis and proliferation to promote mammary epithelial cell transformation," *Nature Structural and Molecular Biology*, vol. 19, no. 2, pp. 220–228, 2012.
- [80] P. E. Czabotar, G. Lessene, A. Strasser, and J. M. Adams, "Control of apoptosis by the BCL-2 protein family: implications for physiology and therapy," *Nature Reviews Molecular Cell Biology*, vol. 15, no. 1, pp. 49–63, 2014.
- [81] A. Shimoni-Sebag, I. Lebenthal-Loinger, L. Zender, and R. Karni, "RRM1 domain of the splicing oncoprotein SRSF1 is required for MEK1-MAPK-ERK activation and cellular transformation," *Carcinogenesis*, vol. 34, no. 11, pp. 2498–2504, 2013.
- [82] H. L. Gautrey and A. J. Tyson-Capper, "Regulation of Mcl-1 by SRSF1 and SRSF5 in cancer cells," *PLoS ONE*, vol. 7, no. 12, Article ID e51497, 2012.
- [83] S. Leu, Y.-M. Lin, C.-H. Wu, and P. Ouyang, "Loss of Pnn expression results in mouse early embryonic lethality and cellular apoptosis through SRSF1-mediated alternative expression of Bcl-xS and ICAD," *Journal of Cell Science*, vol. 125, no. 13, pp. 3164–3172, 2012.
- [84] X. Li, J. Wang, and J. L. Manley, "Loss of splicing factor ASF/SF2 induces G2 cell cycle arrest and apoptosis, but inhibits internucleosomal DNA fragmentation," *Genes and Development*, vol. 19, no. 22, pp. 2705–2714, 2005.
- [85] N. A. Olshavsky, C. E. S. Comstock, M. J. Schiewer et al., "Identification of ASF/SF2 as a critical, allele-specific effector of the cyclin D1b oncogene," *Cancer Research*, vol. 70, no. 10, pp. 3975–3984, 2010.
- [86] F. Wang, H. Wang, H. Wu et al., "TEAD1 controls C2C12 cell proliferation and differentiation and regulates three novel target genes," *Cellular Signalling*, vol. 25, no. 3, pp. 674–681, 2013.
- [87] R. Karni, Y. Hippo, S. W. Lowe, and A. R. Krainer, "The splicing-factor oncoprotein SF2/ASF activates mTORC1," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 105, no. 40, pp. 15323–15327, 2008.

- [88] V. Ben-Hur, P. Denichenko, Z. Siegfried et al., "S6K1 alternative splicing modulates its oncogenic activity and regulates mTORC1," *Cell Reports*, vol. 3, no. 1, pp. 103–115, 2013.
- [89] T. Ezponda, M. J. Pajares, J. Agorreta et al., "The oncoprotein SF2/ASF promotes non-small cell lung cancer survival by enhancing survivin expression," *Clinical Cancer Research*, vol. 16, no. 16, pp. 4113–4125, 2010.
- [90] Y. Fu, B. Huang, Z. Shi et al., "SRSF1 and SRSF9 RNA binding proteins promote Wnt signalling-mediated tumorigenesis by enhancing β -catenin biosynthesis," *EMBO Molecular Medicine*, vol. 5, no. 5, pp. 737–750, 2013.
- [91] C. Ghigna, S. Giordano, H. Shen et al., "Cell motility is controlled by SF2/ASF through alternative splicing of the Ron protooncogene," *Molecular Cell*, vol. 20, no. 6, pp. 881–890, 2005.
- [92] P. Jordan, R. Brazão, M. G. Boavida, C. Gespach, and E. Chastre, "Cloning of a novel human Rac1b splice variant with increased expression in colorectal tumors," *Oncogene*, vol. 18, no. 48, pp. 6835–6839, 1999.
- [93] A. Schnelzer, D. Prechtel, U. Knaus et al., "Rac1 in human breast cancer: overexpression, mutation analysis, and characterization of a new isoform, Rac1b," *Oncogene*, vol. 19, no. 26, pp. 3013–3020, 2000.
- [94] P. Matos, J. G. Collard, and P. Jordan, "Tumor-related alternatively spliced Rac1b is not regulated by Rho-GDP dissociation inhibitors and exhibits selective downstream signaling," *The Journal of Biological Chemistry*, vol. 278, pp. 50442–50448, 2003.
- [95] D. Fiegen, L.-C. Haeusler, L. Blumenstein et al., "Alternative splicing of rac1 generates Rac1b, a self-activating GTPase," *Journal of Biological Chemistry*, vol. 279, no. 6, pp. 4743–4749, 2004.
- [96] V. Gonçalves, P. Matos, and P. Jordan, "Antagonistic SR proteins regulate alternative splicing of tumor-related Rac1b downstream of the PI3-kinase and Wnt pathways," *Human Molecular Genetics*, vol. 18, no. 19, pp. 3696–3707, 2009.
- [97] P. Matos and P. Jordan, "Expression of Rac1b stimulates NF- κ B-mediated cell survival and G1/S progression," *Experimental Cell Research*, vol. 305, no. 2, pp. 292–299, 2005.
- [98] P. Matos and P. Jordan, "Increased Rac1b expression sustains colorectal tumor cell survival," *Molecular Cancer Research*, vol. 6, no. 7, pp. 1178–1184, 2008.
- [99] A. Maimon, M. Mogilevsky, A. Shilo et al., "Mnk2 alternative splicing modulates the p38-MAPK pathway and impacts Ras-induced transformation," *Cell Reports*, vol. 7, no. 2, pp. 501–513, 2014.
- [100] F. J. de Miguel, R. D. Sharma, M. J. Pajares, L. M. Montuenga, A. Rubio, and R. Pio, "Identification of alternative splicing events regulated by the oncogenic factor SRSF1 in lung cancer," *Cancer Research*, vol. 74, no. 4, pp. 1105–1115, 2014.
- [101] D. G. Nowak, J. Woolard, E. M. Amin et al., "Expression of pro- and anti-angiogenic isoforms of VEGF is differentially regulated by splicing and growth factors," *Journal of Cell Science*, vol. 121, no. 20, pp. 3487–3495, 2008.
- [102] E. S. Rennel, A. H. R. Varey, A. J. Churchill et al., "VEGF121b, a new member of the VEGF_{xxx}b family of VEGF-A splice isoforms, inhibits neovascularisation and tumour growth in vivo," *British Journal of Cancer*, vol. 101, no. 7, pp. 1183–1193, 2009.
- [103] S. Oltean, M. Gammons, R. Hulse et al., "SRPK1 inhibition in vivo: modulation of VEGF splicing and potential treatment for multiple diseases," *Biochemical Society Transactions*, vol. 40, no. 4, pp. 831–835, 2012.
- [104] M. N. Thobe, D. Gurusamy, P. Pathrose, and S. E. Waltz, "The Ron receptor tyrosine kinase positively regulates angiogenic chemokine production in prostate cancer cells," *Oncogene*, vol. 29, no. 2, pp. 214–226, 2010.
- [105] A. F. Muro, M. Caputi, R. Pariyarath, F. Pagani, E. Buratti, and F. E. Baralle, "Regulation of fibronectin EDA exon alternative splicing: possible role of RNA secondary structure for enhancer display," *Molecular and Cellular Biology*, vol. 19, no. 4, pp. 2657–2671, 1999.
- [106] I. C. Lopez-Mejia, M. de Toledo, F. Della Seta et al., "Tissue-specific and SRSF1-dependent splicing of fibronectin, a matrix protein that controls host cell invasion," *Molecular Biology of the Cell*, vol. 24, no. 20, pp. 3164–3176, 2013.
- [107] L. Adesso, S. Calabretta, F. Barbagallo et al., "Gemcitabine triggers a pro-survival response in pancreatic cancer cells through activation of the MNK2/eIF4E pathway," *Oncogene*, vol. 32, no. 23, pp. 2848–2857, 2013.
- [108] J. C. Shultz, R. W. Goehe, C. S. Murudkar et al., "SRSF1 regulates the alternative splicing of caspase 9 via a novel intronic splicing enhancer affecting the chemotherapeutic sensitivity of non-small cell lung cancer cells," *Molecular Cancer Research*, vol. 9, no. 7, pp. 889–900, 2011.
- [109] J. C. Shultz, R. W. Goehe, D. S. Wijesinghe et al., "Alternative splicing of caspase 9 is modulated by the phosphoinositide 3-kinase/Akt pathway via phosphorylation of SRp30a," *Cancer Research*, vol. 70, no. 22, pp. 9185–9196, 2010.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

