

## LETTERS

# De Bary syndrome and ATP6V0A2-CDG

*European Journal of Human Genetics* (2010) **18**, 526; doi:10.1038/ejhg.2009.218; published online 16 December 2009

We read with interest the nice review of Morava *et al*<sup>1</sup> on the autosomal recessive cutis laxa syndromes. The authors mention the De Bary syndrome and state that the genetic background of the De Bary syndrome has not yet been identified. However, in the paper by Kornak *et al*<sup>2</sup> on impaired glycosylation and cutis laxa caused by mutations in *ATP6V0A2* (ATP6V0A2-CDG according to the novel nomenclature<sup>3,4</sup>), one of the patients (see patient CoFe in Table 1 of Kornak *et al*<sup>2</sup>) shows the full clinical picture of the De Bary syndrome, including cutis laxa, facial dysmorphism, dwarfism, psychomotor retardation, dystonia, congenital hip dysplasia, and corneal dystrophy necessitating repeated corneal transplantation. These data suggest that a subgroup of patients with De Bary syndrome<sup>5</sup> belongs to the spectrum of ATP6V0A2-CDG. Another cause of De Bary syndrome has very recently been identified as mutations in *PYCR1*, coding for a mitochondrial enzyme involved in proline metabolism.<sup>6</sup> Therefore, we recommend a systematic screen for ATP6V0A2-CDG and for mutations in *PYCR1* in patients with De Bary syndrome.

Elisa Leao-Teles<sup>1</sup>, Dulce Quelhas<sup>2</sup>, Laura Vilarinho<sup>2</sup>  
and Jaak Jaeken<sup>3</sup>

<sup>1</sup>Metabolic Unit, Department of Pediatrics, San Joao, Hospital, Porto, Portugal; <sup>2</sup>Department of Clinical Biology, Institute of Medical Genetics Jacinto de Magalhães, Porto, Portugal; <sup>3</sup>Department of Pediatrics, Center for Metabolic Disease, University Hospital Gasthuisberg, Leuven, Belgium  
E-mail: jaak.jaeken@uzleuven.be

- 1 Morava E, Guillard M, Lefeber DJ, Wevers RA: Autosomal recessive cutis laxa syndrome revisited. *Eur J Hum Genet* 2009; **17**: 1099–1110.
- 2 Kornak U, Reyniers E, Dimopoulou A *et al*: Impaired glycosylation and cutis laxa caused by mutations in the vesicular H<sup>+</sup>-ATPase subunit ATP6V0A2. *Nat Genet* 2008; **40**: 32–34.
- 3 Jaeken J, Hennet T, Freeze HH, Matthijs G: On the nomenclature of congenital disorders of glycosylation (CDG). *J Inher Metab Dis* 2008; **31**: 669–672.
- 4 Jaeken J, Hennet T, Matthijs G, Freeze HH: CDG nomenclature: time for a change. *Biochim Biophys Acta* 2009; **1792**: 825–826.
- 5 Kivuva EC, Parker MJ, Cohen MC, Wagner BE, Sobey G: De Bary syndrome: a review of the phenotype. *Clin Dysmorphol* 2008; **17**: 99–107.
- 6 Reversade B, Escande Beillard N, Dimopoulou A *et al*: Mutations in *PYCR1* cause cutis laxa with progeroid features. *Nat Genet* 2009; **41**: 1016–1021.

## Reply to Leao-Teles *et al*

*European Journal of Human Genetics* (2010) **18**, 526; doi:10.1038/ejhg.2009.219; published online 16 December 2009

We appreciate the comments<sup>1</sup> on our review on autosomal recessive cutis laxa syndromes,<sup>2</sup> especially the novel information on the

Portuguese patient CoFe<sup>3</sup> (Table 1, Kornak *et al*), who was diagnosed with a combined N- and O-linked glycosylation disorder and ATP6V0A2-CDG. De Bary syndrome is a challenging diagnosis, and has overlapping features with ARCL type 2, including cutis laxa, short stature, late closure of the fontanel, congenital hip dysplasia, some aspects of facial dysmorphism and psychomotor retardation.<sup>4</sup> Still, dystonic movements and progressive corneal abnormalities are highly suggestive of De Bary syndrome. So far no cobble-stone-like brain dysgenesis has been observed in De Bary syndrome patients. Although the clinical differentiation is not always straightforward, progeroid features, the presence of pergamen-like skin and the absence of subcutaneous fat are distinctive.<sup>2</sup> Interestingly, some of the patients, recently reported to have *PYCR1* mutations, were previously diagnosed with De Bary syndrome,<sup>5</sup> while other De Bary patients were found to not harbour *PYCR1* mutations (unreported data), thus supporting genetic heterogeneity. We would appreciate further description and photo documentation of this unique patient, reported by Leao-Teles, as in our own cohort of six patients with De Bary syndrome and those studied by Kornak *et al* (personal communication) none of the children had either N-linked or O-linked glycosylation abnormalities, nor mutations in the *ATP6V0A2* gene. Eye anomalies are common in N-glycosylation disorders,<sup>6</sup> and strabismus and high myopia have been reported in combined glycosylation disorders.<sup>4</sup> The exceptional observation of corneal abnormalities with a movement disorder in association with ATP6V0A2-CDG indeed widens the range of symptoms evoking glycosylation studies in patients with cutis laxa.

### CONFLICT OF INTEREST

The authors declare no conflict of interest.

Eva Morava<sup>1</sup>, Mailys Guillard<sup>2</sup>, Dirk J Lefeber<sup>2</sup>  
and Ron A Wevers<sup>2</sup>

<sup>1</sup>Department of Paediatrics, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands;  
<sup>2</sup>Laboratory of Paediatrics and Neurology, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands  
E-mail: E.Morava@cukz.umcn.nl

- 1 Leao-Teles E, Quelhas D, Vilarinho L, Jaeken J: De Bary syndrome and ATP6V0A2-CDG. *Eur J Human Genet* 2010; **18**: 526.
- 2 Morava E, Guillard M, Lefeber DJ, Wevers RA: Autosomal recessive cutis laxa syndrome revisited. *Eur J Hum Genet* 2009; **17**: 1099–1110.
- 3 Kornak U, Reyniers E, Dimopoulou A *et al*: Impaired glycosylation and cutis laxa caused by mutations in the vesicular H<sup>+</sup>-ATPase subunit ATP6V0A2. *Nat Genet* 2008; **40**: 32–34.
- 4 Huchtagowder V, Morava E, Kornak U *et al*: Loss-of-function mutations in ATP6V0A2 impair vesicular trafficking, tropoelastin secretion and cell survival. *Hum Mol Genet* 2009; **18**: 2149–2165.
- 5 Reversade B, Escande-Beillard N, Dimopoulou A *et al*: Mutations in *PYCR1* cause cutis laxa with progeroid features. *Nat Genet* 2009; **41**: 1016–1021.
- 6 Morava E, Wosik HN, Sykut-Cegielska J *et al*: Ophthalmological abnormalities in children with congenital disorders of glycosylation type I. *Br J Ophthalmol* 2009; **93**: 350–354.